

## Systematic review report for Question 1b

### Primary PICO question 1b

For women who are positive for HPV types other than 16 or 18 and have negative or pLSIL/dLSIL reflex liquid based cytology (intermediate risk), what is the safety and effectiveness of repeating HPV testing in 12 and 24 months compared to repeating HPV test at 12 months only before returning to 5 yearly screening?

Population	Study design	Intervention	Control	Outcome
Women who are positive for HPV types other than 16 or 18 and have negative or pLSIL/dLSIL (ASC-US/LSIL - Bethesda) liquid based cytology (intermediate risk)	Randomised or pseudo-randomized controlled trial	Repeat HPV test in 12 and 24 months; Colposcopy and reflex LBC if positive or if both negative discharge back to screening	Repeat HPV test in 12 months; Colposcopy if positive and if negative discharge to screening	Cervical cancer mortality Cervical cancer diagnosis Precancerous high grade lesion detection

ASC-US = Atypical squamous cells, undetermined significance (Bethesda 2001); dLSIL = definite LSIL; LBC = liquid-based cytology; LSIL = Low-grade squamous intraepithelial lesion; pLSIL = possible LSIL

Randomised and pseudo-randomised controlled trials directly address the primary PICO question 1b. In the event that no relevant randomised or pseudo-randomised controlled trials were identified an indirect approach was planned with a secondary PICO question focussing on benchmarking ie examining the risk of high grade lesions in women who are positive for hr-HPV types other than 16 or 18 with normal cytology, and comparing these with current risk thresholds for 12-month follow-up or later colposcopy. Definite LSIL cytology was chosen as the benchmark for the level of risk for 12-month follow-up as 12-month follow-up is recommended for definite LSIL cytology in the current guidelines.

### Secondary PICO question 1b

For women undergoing routine cervical screening what is the risk of **subsequent** CIN3+ for women who are positive for HPV oncogenic types other than 16 and 18 and have **negative** cytology compared with women who have pLSIL/dLSIL regardless of HPV status or who have pLSIL/dLSIL and are positive for HPV oncogenic types other than 16 and 18?

Population	Study design	Exposure	Comparator	Outcome
Women undergoing routine cervical screening	Longitudinal prognostic cohort studies	Positive for HPV oncogenic types other than 16 and 18 and have negative cytology	pLSIL (ASC-US) /dLSIL (LSIL - Bethesda) or positive for HPV oncogenic types other than 16/18 and have pLSIL (ASC-US) /dLSIL (LSIL - Bethesda) cytology	CIN3+ CIN2+

ASC-US = Atypical squamous cells, undetermined significance (Bethesda 2001); CIN 3+ = cervical intraepithelial neoplasia grade 3 or worse; CIN 2+ = cervical intraepithelial neoplasia grade 2 or worse; dLSIL = definite LSIL; LSIL = Low-grade squamous intraepithelial lesion; pLSIL = possible LSIL

## 1. Methods

### 1.1. Search for existing relevant guidelines that could be adopted

Relevant recent (2005 onwards) guidelines were identified by scanning the citations identified by the literature search and searching the National Guideline Clearinghouse (<http://guideline.gov/>) and the Guidelines Resource Centre ([www.cancerview.ca](http://www.cancerview.ca)).

To be considered for adoption guidelines had to be directly relevant, based on systematic reviews of the evidence and meet the pre-specified criteria of scores of greater or equal to 70% for the domains rigour of development, clarity of presentation and editorial independence of the AGREE II instrument (<http://www.agreetrust.org/resource-centre/agree-ii/>).

### 1.2 Literature search

In 2013 systematic reviews were undertaken by the NHMRC Clinical Trials Centre (CTC) to provide the evidence base for the assessment by the Medical Services Advisory Committee (MSAC) of cervical screening strategies for the renewal of the National Cervical Screening Program ([http://www.cancerscreening.gov.au/internet/screening/publishing.nsf/Content/E6A211A6FFC29E2CC A257CED007FB678/\\$File/Review%20of%20Evidence%20notated%2013.06.14.pdf](http://www.cancerscreening.gov.au/internet/screening/publishing.nsf/Content/E6A211A6FFC29E2CC A257CED007FB678/$File/Review%20of%20Evidence%20notated%2013.06.14.pdf)).

One of the CTC systematic reviews addressed the question,

*What is the comparative safety and effectiveness of undertaking a colposcopy immediately in comparison to delaying the test in women who have pLSIL/dLSIL cytology and a positive HPV test? (MSAC Question 3.3).*

This CTC review would include any studies relevant to primary question 1b. Given that this review was based on searches using detailed and comprehensive search strategies of multiple databases up until December 2012 and the application of precise and transparent inclusion criteria, it was used as the source of evidence up until the end of 2011. To identify subsequent relevant publications, Medline, the Cochrane Central Register of Controlled Trials (CENTRAL), Embase, PreMedline, Database of Abstracts of Reviews of Effects (DARE) and Health Technology Assessments (HTA) databases were searched for articles published from 2012 until 31<sup>st</sup> August 2015, using text terms and, where available, database-specific subject headings. In these databases searches for cervical cytology were combined with searches for HPV and colposcopy, and where possible, database-specific filters for identifying randomized controlled trials. To identify studies which considered Aboriginal and Torres Strait Islander (ATSI) peoples these searches were then coupled with search terms for ATSI peoples. A complete list of the terms used for search strategies are included as Appendix A. The Cochrane systematic review database was also searched for relevant systematic reviews or meta-analyses using the terms, HPV and colposcopy, and abstracts from the 2015 EUROGIN conference were scanned for relevant studies using the term "colpos". Reference lists of relevant articles and guidelines were checked for additional potentially relevant articles.

To identify publications that addressed the secondary PICO question the Medline, PreMedline, Embase, and CENTRAL databases were searched for articles published from 2004 until 31<sup>st</sup> August 2015, using the same search strategies with the randomised controlled trial filters removed, and the

results of the DARE, HTA, the Cochrane systematic review database and the EUROGIN 2015 abstracts searches undertaken for the primary PICO question were reassessed against the broader inclusion criteria of the secondary PICO. A complete list of the terms used for these search strategies are included in Appendix A.

### 1.1. Inclusion criteria

#### Inclusion criteria for primary PICO question 1b

Selection criteria	Inclusion criteria	Exclusion criteria
Population	Women undergoing screening who are positive for oncogenic HPV types other than 16 or 18 and have Negative or pLSIL (ASC-US – Bethesda 2001) or dLSIL (LSIL - Bethesda) cytology	Conducted in referred population and do not specifically exclude women undergoing follow-up following treatment For HPV test used <ul style="list-style-type: none"> <li>• In-situ hybridization or</li> <li>• p16 immunostaining or</li> <li>• Tests of viral load or</li> <li>• Hybrid Capture I or</li> </ul> HPV test detected oncogenic HPV types other than those detected by the Cobas test
Study type	Screening intervention	
Study design	Randomised or pseudo-randomised controlled trials or Systematic review/meta-analyses thereof	Reviews, editorials Conference proceedings other than 2015 EUROGIN conference proceedings
Intervention	Colposcopy dependent on subsequent repeat oncogenic HPV testing at 12 (+/- 6 months) and 24 (+/- 6 months) with discharge to screening if HPV negative on both tests	Repeat HPV test delayed to 24 months <sup>^</sup>
Comparison	Colposcopy dependent on repeat oncogenic HPV testing at 12 (+/- 6 months) with discharge to screening if HPV negative	
Outcomes	Cervical cancer mortality or Cervical cancer diagnosis or CIN3 <sup>^</sup> diagnosis or CIN2+ diagnosis	
Search period	1 <sup>st</sup> January 2004 – 31 <sup>st</sup> August 2015	
Language	English	

ASC-US = Atypical squamous cells, undetermined significance (Bethesda 2001); CIN2+ = cervical intraepithelial neoplasia grade 2 or worse; CIN3+ = cervical intraepithelial neoplasia grade 3 or worse; dLSIL = definite LSIL; LSIL = Low-grade squamous intraepithelial lesion; pLSIL = possible LSIL

<sup>^</sup> MSAC states “Cervical cancer is rare in screened populations and therefore trials designed to detect differences in incidence and mortality of cervical cancer in these populations are unfeasible. Reduced CIN3+ is considered as an acceptable proxy outcome of trials evaluating new preventative strategies (Arbyn et al. 2009).”

<sup>^</sup>The ARTISTIC study modelling showed that the risk of CIN2+ is higher in women who are referred at 24 months instead of 12 months and is not safe for women positive for HPV at baseline

## Inclusion criteria for secondary PICO question 1a

Selection criteria	Inclusion criteria	Exclusion criteria
Population	Women undergoing routine screening	Conducted in referred population and do not specifically exclude women undergoing follow-up following treatment
Study type	Prognostic	
Study design	Cohort: Longitudinal	Case-control studies in which absolute risk cannot be calculated Reviews, editorials Conference proceedings other than 2015 EUROGIN conference proceedings Cross-sectional cohort studies – no follow-up
Exposure	Positive for oncogenic HPV types other than 16 or 18 with negative cytology	For HPV test used <ul style="list-style-type: none"> <li>• In-situ hybridization or</li> <li>• p16 immunostaining or</li> <li>• Tests of viral load or</li> <li>• Hybrid Capture I or</li> </ul> HPV test detected oncogenic HPV types other than those detected by the Cobas test
Comparator	dLSIL (LSIL - Bethesda) cytology regardless of HPV status	Exposed and comparators assessed in a different age groups  No exit colposcopy for all participants and follow-up differed for exposed and comparators ie reference standard of colposcopy and/or biopsy not systematically applied to both exposed and comparator  Assessed only a subgroup of comparators eg HPV+ve LSIL
Outcomes	Cervical cancer diagnosis or CIN3+ diagnosis or CIN2+ diagnosis	
Search period	1 <sup>st</sup> January 2004 – 31 <sup>st</sup> August 2015	
Language	English	

CIN2+ = cervical intraepithelial neoplasia grade 2 or worse; CIN3+ = cervical intraepithelial neoplasia grade 3 or worse; dLSIL = definite LSIL; LSIL = Low-grade squamous intraepithelial lesion;

## 2. Results

### 2.1. Results of Guidelines search

The American Cancer Society, American Society for Colposcopy and Cervical Pathology, and American Society for Clinical Pathology 2012 *Screening guidelines for the prevention and early detection of cervical cancer* contained potentially relevant recommendations regarding cervical screening protocols using partial HPV genotyping. These recommendations were included in the American Society for Colposcopy and Cervical Pathology's 2012 *Updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors* and the American College of Obstetricians and Gynecologists' 2012 *Screening for cervical cancer guidelines*. They were not adopted as they considered follow-up using co-testing not HPV testing and thus did not specifically address the primary PICO question.

## 2.2. Results of Literature Search

### Primary PICO


The CTC systematic review did not include any studies that met the inclusion criteria for primary PICO question 1a. Figure 1 outlines the steps undertaken to identify relevant published from 2012 onwards. Searches of the Medline, PreMedline, Embase and CENTRAL databases identified 156 citations, DARE and HTA databases another 22 citations, the Cochrane database of systematic reviews 73 citations and EUROGIN abstracts, 16 citations; a total of 134 citations. Titles and abstracts were examined and 15 articles were retrieved for a more detailed evaluation.

None met the inclusion criteria for the primary PICO thus **no studies were found that directly addressed the primary PICO question.**

There were no studies of Aboriginal and/or Torres Strait Islander women that met the inclusion criteria.

The retrieved articles that were not included and the reason for their exclusion are documented in Appendix C. The main reasons for exclusion were an absence of outcomes specifically for women with negative, pLSIL or dLSIL cytology who were positive for oncogenic HPV types other than 16 or 18, did not identify for women with negative, pLSIL or dLSIL cytology who were positive for oncogenic HPV types other than 16 or 18 or were not randomised or pseudo-randomised controlled trials.

Database or Source	Number of Citations	Number of Articles Collected	Number of Articles Included	ATSI filter results
Medline and PreMedline and CENTRAL	116	9	0	NA
Embase	40	4	0	NA
Cochrane database of systematic reviews	73	0	0	NA
HTA and DARE	22	2	0	NA
EUROGIN	16	NA	0	NA
<b>TOTAL</b>	<b>267</b>	<b>15</b>	<b>0</b>	<b>NA</b>



**Figure 1. Process of inclusion and exclusion of studies for primary PICO questions**


### Secondary PICO

Searches were then broadened as pre-planned to identify studies that might address the secondary PICO question. Figure 2 outlines the steps undertaken to identify relevant articles for the systematic review. Searches of the Medline, Embase, PreMedline and CENTRAL databases with the RCT filter removed identified 2,551 citations. When combined with those identified by HTA and DARE database and 2015 EUROGIN abstracts searches, a total of 2,671 citations were identified. Titles and abstracts were examined and 413 articles were retrieved for a more detailed evaluation. An additional 9 potential citations were identified from the reference list of retrieved articles resulting in a total of 422 retrieved articles.

**No studies** were found that met the inclusion criteria for the secondary PICO question systematic review. There were no studies of Aboriginal and/or Torres Strait Islander women that met the inclusion criteria.

The retrieved articles that were not included and the reason for their exclusion are documented in Appendix C. The main reason for exclusion was an absence of outcomes for women positive for oncogenic HPV types other than 16 and /or 18 with negative cytology.

Database or Source	Number of Citations	Number of Articles Collected	Number of Articles Included	ATSI filter results
Medline and PreMedline and CENTRAL	1,550	9 + 354	0	NA
Embase	1,001	48	0	NA
Cochrane database of systematic reviews	73	0	0	NA
HTA and DARE	22	2	0	NA
EUROGIN	16	NA	0	NA
snowballing	NA	9	0	NA
<b>TOTAL</b>	<b>2,662</b>	<b>422</b>	<b>0</b>	<b>NA</b>



**Figure 2. Process of inclusion and exclusion of studies**



### 3. Appendices

#### Appendix A: Search strategies used

For Medline, Premedline and Cochrane Central Register of Controlled trials databases (Ovid):

#	Searches
1	HPV.mp.
2	hr\$HPV.mp.
3	papillomavirus.mp.
4	exp Papillomavirus Infections/
5	exp Papillomaviridae/
6	exp DNA Probes, HPV/
7	1 or 2 or 3 or 4 or 5 or 6
8	hybrid capture.mp.
9	HC2.mp.
10	HCII.mp.
11	(hybrid adj5 capture).mp.
12	realtime.mp.
13	cervista.mp.
14	amplicor.mp.
15	cobas 4800.mp.
16	linear array.mp.
17	(linear adj3 array).mp.
18	papillocheck.mp.
19	8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18
20	exp Polymerase Chain Reaction/
21	pcr.mp.
22	20 or 21
23	7 or 19 or 22
24	exp Vaginal Smears/
25	((cervi* or vagina*) and (smear* or screening* or test*)).tw.
26	(pap adj5 smear).tw.
27	papanicolaou.tw.
28	LBC.mp.
29	cytolog\$.mp.
30	exp Uterine Cervical Dysplasia/
31	cervical dysplasia.mp.
32	(cervi* adj5 dysplasia).mp.
33	(dyskaryosis or dyskariosis).mp.
34	ASCUS.mp.
35	ASC US.mp.
36	ASC\$US.mp.
37	ASC R.mp.

38	(atypical squamous adj4 cervi\$.mp.
39	atypical endocervical.mp.
40	atypical gland\$.mp.
41	AGUS.mp.
42	((borderline or low-grade) adj3 abnormal\$.mp.
43	((borderline or low-grade) adj3 cytology).mp.
44	SIL.mp.
45	LSIL.mp.
46	L-SIL.mp.
47	LGSIL\$.mp.
48	pLSIL\$.mp.
49	dLSIL\$.mp.
50	low-grade squamous intraepithelial.mp.
51	possible low-grade squamous intraepithelial.mp.
52	definite low-grade squamous intraepithelial.mp.
53	24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52
54	exp Colposcopy/
55	colposcop\$.mp.
56	54 or 55
57	Surveillance.mp.
58	56 or 57
59	23 and 53 and 58
60	randomized controlled trial.pt.
61	controlled clinical trial.pt.
62	placebo.ab.
63	randomi?ed.ab.
64	randomly.ab.
65	trial.ab.
66	groups.ab.
67	60 or 61 or 62 or 63 or 64 or 65 or 66
68	59 and 67
69	limit 68 to english language

Used the Cochrane sensitivity maximizing filter for identifying randomized controlled trials (<http://handbook.cochrane.org>, accessed 12/09/2015)

#### ATSI search terms used

#	Searches
1	((exp Australia/ OR Australia\$.ti,ab) AND (Oceanic ancestry group/ OR aborigin\$.ti,ab. OR indigenous.mp.)) OR torres strait\$ islander\$.ti,ab

From the Lowitja Institute at <http://www.lowitja.org.au/litsearch-background-information> accessed 30/09/2013)

For Embase database (Embase):

#	Searches
1.	'hpv 16'/exp OR 'hpv 16'
2.	'hpv18'/exp OR 'hpv18'
3.	'hpv31'/exp OR 'hpv31'
4.	'hpv33'/exp OR 'hpv33'
5.	'hpv35'/exp OR 'hpv35'
6.	'hpv45'/exp OR 'hpv45'
7.	'hpv 52'/exp OR 'hpv 52'
8.	'hpv 58'/exp OR 'hpv 58'
9.	'hpv'/exp OR 'hpv'
10.	'human papillomavirus'/exp OR 'human papillomavirus'
11.	'human papillomavirus test'
12.	hr*hpv
13.	#1 OR #2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9 OR #10 OR #11 OR #12
15.	'vagina smear'/exp OR 'vagina smear'
16.	'uterine cervix cytology'/exp OR 'uterine cervix cytology'
17.	'uterine cervix dysplasia'/exp OR 'uterine cervix dysplasia'
18.	cervi* OR vagina* AND (smear* OR screen* OR test*)
19.	pap* NEAR/5 smear*
20.	papanicolaou
21.	'papanicolaou test'/exp OR 'papanicolaou test'
22.	lbc
23.	cytolog*
24.	cervi* NEAR/5 dysplasia
25.	dyskaryosis OR dyskariosis
26.	ascus OR 'asc us' OR 'asc r'
27.	'asc-us' OR 'asc-r'
28.	atypical AND squamous NEAR/4 cervi*
29.	'atypical endocervical'
30.	atypical NEXT/1 gland*
31.	agus
32.	(borderline OR 'low-grade') NEAR/3 abnormal*
33.	(borderline OR 'low-grade') NEAR/3 cytology
35.	sil* OR lsil* OR 'l-sil' OR lgsil* OR plsil* OR dlsil* OR 'p/dlsil'

36.	'low-grade squamous intraepithelial'
37.	#15 OR #16 OR #17 OR #18 OR #19 OR #20 OR #21 OR #22 OR #23 OR #24 OR #25 OR #26 OR #27 OR #28 OR #29 OR #30 OR #31 OR #32 OR #33 OR #35 OR #36
38.	'colposcopy'/exp
39.	colposcop*
40.	surveillance
41.	#38 OR #39 OR #40
42.	rct
43.	'randomized controlled trial'/de
44.	'randomised controlled trial' OR 'randomized controlled trial' OR 'randomised controlled trials' OR 'randomized controlled trials'
45.	'random allocation' OR 'randomly allocated'
46.	'randomization'/de
47.	allocated NEAR/2 random
48.	'double blind procedure'/de
49.	'single blind procedure'/de
50.	single NEXT/1 blind*
51.	double NEXT/1 blind*
52.	(treble OR triple) NEXT/1 blind*
53.	placebo
54.	'placebo'/de
55.	'prospective study'/de
56.	'crossover procedure'/de
57.	'clinical trial'/de
58.	#42 OR #43 OR #44 OR #45 OR #46 OR #47 OR #48 OR #49 OR #50 OR #51 OR #52 OR #53 OR #54 OR #55 OR #56 OR #57
59.	'case study'/de
60.	'case report'
61.	'abstract report'/de
62.	'letter'/de
63.	#59 OR #60 OR #61 OR #62
64.	#58 NOT #63
65.	#13 AND #37 AND #41 AND #64
66.	#13 AND #37 AND #41 AND #64 AND [2013-2015]/py
67.	#13 AND #37 AND #41 AND #64 AND [2013-2015]/py AND [english]/lim
68.	#13 AND #37 AND #41 AND #64 AND [2013-2015]/py AND [english]/lim AND [embase]/lim

69.	#13 AND #37 AND #41 AND #64 AND [2013-2015]/py AND [english]/lim AND [embase]/lim AND [medline]/lim
70.	#67 NOT #69

Used the SIGN filter for identifying randomized controlled trials ([www.sign.ac.uk/methodology/filters.html#systematic](http://www.sign.ac.uk/methodology/filters.html#systematic) accessed 20/02/2013)

#### ATSI search terms used

#	Searches
1	'australia'/exp OR australia*:ab,ti
2	'aborigine'/exp OR aborigin*:ab,ti OR indigenous:de,ab,ti
3	'torres strait islander':ab,ti OR 'torres strait islanders':ab,ti
4	#1 AND #2 OR #3

For Health Technology Assessments (HTA) and Database of Abstracts of Reviews of Effects (DARE) databases (Ovid):

#	Searches
1	HPV.mp.
2	hr\$HPV.mp.
3	papillomavirus.mp.
4	exp Papillomavirus Infections/
5	exp Papillomaviridae/
6	exp DNA Probes, HPV/
7	1 or 2 or 3 or 4 or 5 or 6
8	hybrid capture.mp.
9	HC2.mp.
10	HCII.mp.
11	(hybrid adj5 capture).mp.
12	realtime.mp.
13	cervista.mp.
14	amplacor.mp.
15	cobas 4800.mp.
16	linear array.mp.
17	(linear adj3 array).mp.
18	papillocheck.mp.
19	8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18
20	exp Polymerase Chain Reaction/
21	pcr.mp.
22	20 or 21
23	7 or 19 or 22
24	exp Vaginal Smears/
25	((cervi* or vagina*) and (smear* or screening* or test*)).tw.

26	(pap adj5 smear).tw.
27	papanicolaou.tw.
28	LBC.mp.
29	cytolog\$.mp.
30	exp Uterine Cervical Dysplasia/
31	cervical dysplasia.mp.
32	(cervi* adj5 dysplasia).mp.
33	(dyskaryosis or dyskariosis).mp.
34	ASCUS.mp.
35	ASC US.mp.
36	ASC\$US.mp.
37	ASC R.mp.
38	(atypical squamous adj4 cervi\$).mp.
39	atypical endocervical.mp.
40	atypical gland\$.mp.
41	AGUS.mp.
42	((borderline or low-grade) adj3 abnormal\$).mp.
43	((borderline or low-grade) adj3 cytology).mp.
44	SIL.mp.
45	LSIL.mp.
46	L-SIL.mp.
47	LGSIL\$.mp.
48	pLSIL\$.mp.
49	dLSIL\$.mp.
50	low-grade squamous intraepithelial.mp.
51	possible low-grade squamous intraepithelial.mp.
52	definite low-grade squamous intraepithelial.mp.
53	24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35 or 37 or 38 or 39 or 40 or 41 or 42 or 43 or 44 or 45 or 46 or 47 or 48 or 49 or 50 or 51 or 52
54	exp Colposcopy/
55	colposcop\$.mp.
56	54 or 55
57	Surveillance.mp.
58	56 or 57
59	23 and 53 and 58
60	limit 59 to english language

For Cochrane Database of Systematic Reviews:

#	Searches
1	colposcop\$.mp.
2	HPV.mp.
3	1 or 2

## Appendix B:

### Level of evidence rating criteria – Risk assessment studies

Level	Study design
I	Meta-analysis or a systematic review of level II studies
II	Prospective cohort studies
III-1	All or none
III-2	Retrospective cohort studies
III-3	Case control studies
IV	Cross-sectional studies or case series

*According to the standards of the National Health and Medical Research Council*

### Relevance of the Evidence

Rating	Relevance
1	Evidence of an effect on patient-relevant clinical outcomes including benefits and harms, quality of life and survival.
2	Evidence of an effect on a surrogate outcome* that has been shown to be predictive of patient-relevant outcomes for the same intervention.
3	Evidence of an effect on proven surrogate outcomes but for a different intervention.
4	Evidence of an effect on proven surrogate outcomes but for a different intervention and population.
5	Evidence confined to unproven surrogate outcomes.

*\*surrogate outcome' refers to reasonable indicators of whether there has been some effect (e.g. blood pressure measurements or levels of serum cholesterol)*

#### Points for considering patient-relevant outcomes:

- i) The goal of decision making in health care is to choose the intervention(s) (which may include doing nothing) that is (are) most likely to deliver the outcomes that patients find desirable.
- ii) Surrogate outcomes (such as blood pressure measurements or levels of serum cholesterol) may be reasonable indicators of whether there has been some effect. However, they should not be the basis for clinical decisions unless they reliably predict an effect on the way the patient feels, otherwise they will not be of interest to the patient or their carers.
- iii) All possible outcomes that are of most interest to patients (particularly harms) should be identified and evaluated.

*Adapted from table 1.10 of: National Health and Medical Research Council. How to use the evidence: assessment and application of scientific evidence. Canberra: NHMRC; 2000.*

<http://www.nhmrc.gov.au/files/nhmrc/file/publications/synopses/cp69.pdf>

## Appendix C:

### Excluded Studies

#### Primary PICO question

Study	Reason for Exclusion
Bergeron 2015	Does not include women with relevant HPV and cytology status
Blatt 2015	Does not include women with relevant HPV and cytology status
Cruickshank 2015	No relevant comparison
Cuzick 2013	Not randomised or pseudo-randomised controlled trial
Del Mistro 2014	Does not include women with relevant HPV and cytology status
Dijkstra 2014	No relevant comparison
Gage 2013	Not randomised or pseudo-randomised controlled trial
Kececioglu 2013	No results for women with relevant HPV and cytology status
Kitchener 2014	No results for women with relevant HPV and cytology status
McIntosh 2012	Review
Nakamura 2015	Not randomised or pseudo-randomised controlled trial
Reid 2015	No results for women with relevant HPV and cytology status
Ronco 2014	No results for women with relevant HPV and cytology status
Van Le 2015	Summary of a published study
Wright 2013	Conference proceedings 2013

#### Secondary PICO question

Study	Reason for Exclusion
Adamopoulou 2009	No results for women with relevant HPV and cytology status
Agodi 2009	No results for women with relevant HPV and cytology status
Agorastas 2005	No results for women with relevant HPV and cytology status
Agorastos 2015	No relevant comparators
Aksu 2011	Conference proceedings
Alameda 2014	Conference proceedings
Alamonte 2007	No results for women with relevant HPV and cytology status
Allia 2015	No results for women with relevant HPV and cytology status
Annappa 2015	Conference proceedings
Antonishyn 2008	No results for women with relevant HPV and cytology status
Antonishyn 2009	No results for women with relevant HPV and cytology status
Arbyn 2004	No results for women with relevant HPV and cytology status
Arbyn 2009	No results for women with relevant HPV and cytology status
Arbyn 2012	No results for women with relevant HPV and cytology status
Atkins 2006	Does not assess women with relevant HPV and cytology status
Ault 2011	Participants women with AIS
Badano 2011	No results for women with relevant HPV and cytology status
Bais 2005	No results for women with relevant HPV and cytology status



Balbi 2012	No results for women with relevant HPV and cytology status
Bapir 2013	Conference proceedings
Barcelos 2011	No results for women with relevant HPV and cytology status
Barzon 2011	No results for women with relevant HPV and cytology status
Basu 2011	Cervical cancer specimens only
Bekkers 2004	No results for women with relevant HPV and cytology status
Belinson 2010	No results for women with relevant HPV and cytology status
Belinson 2011	No results for women with relevant HPV and cytology status
Bello 2009	No results for women with relevant HPV and cytology status
Benevolo 2011	No results for women with relevant HPV and cytology status
Benevolo 2011b	No results for women with relevant HPV and cytology status
Benoy 2011	No results for women with relevant HPV and cytology status
Bergeron 2015	Does not include women with relevant HPV and cytology status
Berkhof 2006	No results for women with relevant HPV and cytology status
Bhatla 2007	No results for women with relevant HPV and cytology status
Bhatla 2008	No results for women with relevant HPV and cytology status
Bian 2013	Unclear as to whether results for women with relevant HPV and cytology status
Bjerre 2008	No results for women with relevant HPV and cytology status
Blatt 2015	Does not include women with relevant HPV and cytology status
Boardman 2006	Does not include women with relevant HPV and cytology status
Boldrini 2014	No results for women with relevant HPV and cytology status
Bollmann 2006	Post treatment follow-up population
Bourgain 2011	Conference proceedings
Bratti 2004	No results for women with relevant HPV and cytology status
Brink 2006	No results for women with relevant HPV and cytology status
Bulk 2007	No comparable comparative data
Bulkmans 2007	No results for women with relevant HPV and cytology status
Bulkmans 2007b	No results for women with relevant HPV and cytology status
Campari 2015	No results for women with relevant HPV and cytology status
Canda 2009	No results for women with relevant HPV and cytology status
Carozzi 2005	No results for women with relevant HPV and cytology status
Carozzi 2006	No results for women with relevant HPV and cytology status
Carozzi 2013	No results for women with relevant HPV and cytology status
Carozzi 2013b	No results for women with relevant HPV and cytology status
Carter 2010	No results for women with relevant HPV and cytology status
Castle 2004	Commentary
Castle 2007	Participants women with CIN2 histology on colposcopy
Castle 2008	No results for women with relevant HPV and cytology status
Castle 2009	No results for women with relevant HPV and cytology status
Castle 2009 short	No results for women with relevant HPV and cytology status
Castle 2009c	No outcomes of interest
Castle 2009d	No results for women with relevant HPV and cytology status

Castle 2010	Does not include women with relevant HPV and cytology status
Castle 2011a	No results for women with relevant HPV and cytology status
Castle 2011b	No results for women with relevant HPV and cytology status
Castle 2011c	Participants women who have CIN1 or less
Castle 2012	No results for women with relevant HPV and cytology status
Castle 2015	No results for women with relevant HPV and cytology status
Castle 2015 Comarison	No results for women with relevant HPV and cytology status
Castle 2015 Reliability	No results for women with relevant HPV and cytology status
Cattani 2009	No results for women with relevant HPV and cytology status
Chao 2008	No results for women with relevant HPV and cytology status
Chao 2010	No results for women with relevant HPV and cytology status
Chen 2006	No results for women with relevant HPV and cytology status
Chen 2015	No results for women with relevant HPV and cytology status
Chevarie-Davis 2013	No results for women with relevant HPV and cytology status
Chiang 2013	No results for women with relevant HPV and cytology status
Chivukula 2006	Does not include women with relevant HPV and cytology status
Cho 2015	No results for women with relevant HPV and cytology status
Chranioti 2012	No results for women with relevant HPV and cytology status
Cilingir 2013	Colposcopy referral population and unclear if follow-up patients excluded
Ciotto 2004	No results for women with relevant HPV and cytology status
Clad	Includes post treatment follow-up population
Clarke 2014	Conference proceedings
Cobo 2009	No results for women with relevant HPV and cytology status
Confortini 2010	No results for women with relevant HPV and cytology status
Corneanu 2011	No results for women with relevant HPV and cytology status
Correa 2012	No results for women with relevant HPV and cytology status
Cottier 2009	No results for women with relevant HPV and cytology status
Cotton 2010	No results for women with relevant HPV and cytology status
Cox 2013	No relevant comparator
Cruickshank 2014	Results for only a subgroup of women with relevant HPV and cytology status
Crum 2004	Participants women with CIN2/CIN3
Cuschieri 2014	No results for women with relevant HPV and cytology status
Cuzick 2008	No results for women with relevant HPV and cytology status
Cuzick 2010	No results for women with relevant HPV and cytology status
Cuzick 2013	No results for women with relevant HPV and cytology status
Dalla Palma 2008	Participants women with CIN histology
Dalstein 2004	No results for women with relevant HPV and cytology status
Dane 2009	No results for women with relevant HPV and cytology status
Daponte 2006	No results for women with relevant HPV and cytology status
de Vuyst 2005	No results for women with relevant HPV and cytology status
Del Mistro 2010	No results for women with relevant HPV and cytology status
Del Mistro 2014	Does not include women with relevant HPV and cytology status
Denny 2010	No results for women with relevant HPV and cytology status

Depuydt 2011	No results for women with relevant HPV and cytology status
Desteli 2004	No results for women with relevant HPV and cytology status
Diaz Montes 2007	Not relevant population - AGUS or AGC cytology
Dijkstra 2013	No results for women with relevant HPV and cytology status
Dillner 2008	No results for women with relevant HPV and cytology status
Dillner 2011	No results for women with relevant HPV and cytology status
Dinc 2010	No results for women with relevant HPV and cytology status
Ding 2011	Participants women treated for CIN2 histology
Ding 2014	No results for women with relevant HPV and cytology status
Discacciati 2014	Not a cervical screening population - women with CIN2
Dockter 2009	Includes follow-up population
Du 2011	No results for women with relevant HPV and cytology status
Dufresne 2011	No results for women with relevant HPV and cytology status
Einstein 2010	No results for women with relevant HPV and cytology status
Einstein 2011	No results for women with relevant HPV and cytology status
Elfgren 2005	No results for women with relevant HPV and cytology status
Eltoum 2005	No results for women with relevant HPV and cytology status
Filho 2005	No results for women with relevant HPV and cytology status
Fokom-Domgue 2015	No results for women with relevant HPV and cytology status
Froberg 2008	No results for women with relevant HPV and cytology status
Gage 2010	ASCUS/LSIL: <=CIN1 on immediate colposcopy
Gage 2013	No results for women with relevant HPV and cytology status
Gage 2014	No results for women with relevant HPV and cytology status
Geng 2011	Conference proceedings
Geraets 2013	No results for women with relevant HPV and cytology status
Gilani 2013	No results for women with relevant HPV and cytology status
Giorgi Rossi 2010	No results for women with relevant HPV and cytology status
Giorgi Rossi 2011	No results for women with relevant HPV and cytology status
Giorgi Rossi 2013	No results for women with relevant HPV and cytology status
Giovannelli 2005	No results for women with relevant HPV and cytology status
Girianelli 2006	No results for women with relevant HPV and cytology status
Gold 2013	No results for women with relevant HPV and cytology status
Gonzalez- Bosquet 2006	No results for women with relevant HPV and cytology status
Gonzalez-Bosquet 2008	No results for women with relevant HPV and cytology status
Gonzalez-Losa 2004	No results for women with relevant HPV and cytology status
Goodrich 2014	No relevant comparator
Gravitt 2010	No results for women with relevant HPV and cytology status
Grisaru 2008	Not a cervical screening population - included women undergoing post treatment follow-up
Guo 2014	No results for women with relevant HPV and cytology status
Gurumurthy 2014	Participants women with CIN1
Gutierrez-Delgado 2014	Conference proceedings
Haidopoulos 2009	No results for women with relevant HPV and cytology status
Haldorsen 2015	No results for women with relevant HPV and cytology status
Halfon 2010	No results for women with relevant HPV and cytology status

Halfon 2013	No results for women with relevant HPV and cytology status
Hariri 2015	Population women with CIN2
Hariri 2015b	Population women with CIN2
HooKim 2014	Does not include women with relevant HPV and cytology status
Hopenhayn 2014	Not a cervical screening population - women with cervical cancer
Hou 2012	No results for women with relevant HPV and cytology status
Hoyer 2005	No results for women with relevant HPV and cytology status
Hu 2011	No results for women with relevant HPV and cytology status
Huang 2006	No results for women with relevant HPV and cytology status
Huang 2009	No results for women with relevant HPV and cytology status
Hughes 2005	No results for women with relevant HPV and cytology status
Huh 2015	Clinical guidelines
Ibanez 2014	No results for women with relevant HPV and cytology status
Iftner 2015	No results for women with relevant HPV and cytology status
Ikenberg 2013	No results for women with relevant HPV and cytology status
Inoue 2007	No results for women with relevant HPV and cytology status
Jakobsson 2012	Participants women with recurrent cytological abnormalities
Jariene 2012	No results for women with relevant HPV and cytology status
Jentschke 2013	No results for women with relevant HPV and cytology status
Jeronimo 2015	No results for women with relevant HPV and cytology status
Jiang 2011	No results for women with relevant HPV and cytology status
Johnson 2008	No results for women with relevant HPV and cytology status
Jones 2011	Editorial
Jovanovic 2012	No results for women with relevant HPV and cytology status
Kakti 2011	No results for women with relevant HPV and cytology status
Kakti 2013 Benchmarking	No results for women with relevant HPV and cytology status
Kakti 2013 CIN1	Participants women with HPV infection or abnormal cytology and histology <CIN2
Kares 2014	Conference proceedings
Katki 2013 LSIL	No results for women with relevant HPV and cytology status
Katki 2013d neg	No results for women with relevant HPV and cytology status
Katki 2013f ASCUS	No results for women with relevant HPV and cytology status
Kececioglu 2013	No results for women with relevant HPV and cytology status
Keegan 2014	Participants women with recurrent cytological abnormalities
Kelly 2011	No results for women with relevant HPV and cytology status
Khan 2005	Participants included women with prior history of HSIL/CIN2+ diagnosis
Kim 2013	No results for women with relevant HPV and cytology status
Kitchener 2008	Post treatment population
Kitchener 2011	No results for women with relevant HPV and cytology status
Kitchener 2014	No results for women with relevant HPV and cytology status
Ko 2011	No results for women with relevant HPV and cytology status
Kocken 2012	No results for women with relevant HPV and cytology status
Kotaniemi-Talonen 2005	No results for women with relevant HPV and cytology status
Kotaniemi-Talonen 2008	No results for women with relevant HPV and cytology status
Kovacic 2008	No results for women with relevant HPV and cytology status

Krambeck 2008	No results for women with relevant HPV and cytology status
Kroupis 2007	No results for women with relevant HPV and cytology status
Kulasingam 2006	No results for women with relevant HPV and cytology status
Lai 2008	No relevant comparator
Lazcano-Ponce 2010	No results for women with relevant HPV and cytology status
Le Donne 2013	No results for women with relevant HPV and cytology status
Lee 2004	No results for women with relevant HPV and cytology status
Lee 2006	No results for women with relevant HPV and cytology status
Lee 2009	No results for women with relevant HPV and cytology status
Lee 2015	No comparator
Leeson 2014	Discussion article
Leinonen 2013	No results for women with relevant HPV and cytology status
Li 2015	No results for women with relevant HPV and cytology status
Liao 2014	No results for women with relevant HPV and cytology status
Lin 2011	No results for women with relevant HPV and cytology status
Liou 2015	No results for women with relevant HPV and cytology status
Litjens 2013	No results for women with relevant HPV and cytology status
Litjens 2014	Participants women with CIN3 who were previously CIN1
Littell 2011	No results for women with relevant HPV and cytology status
Liu 2014	No results for women with relevant HPV and cytology status
Liverani 2012	No results for women with relevant HPV and cytology status
Loiacono 2012	Conference proceedings
Louvanto 2010	No results for women with relevant HPV and cytology status
Louwers 2010	No results for women with relevant HPV and cytology status
Luttmer 2015	No results for women with relevant HPV and cytology status
Maenpaa 2014	Conference proceedings
Massad 2013	Consensus guidelines
Massad 2014	Review
Maucourt-Boulch 2009	No results for women with relevant HPV and cytology status
Mayrand 2007	No results for women with relevant HPV and cytology status
McDonald 2014	No results for women with relevant HPV and cytology status
McIntosh 2012	Review
McKenna 2014	No results for women with relevant HPV and cytology status
Medeiros 2005	No results for women with relevant HPV and cytology status
Mendoza 2011	No results for women with relevant HPV and cytology status
Mentzelopoulou 2014	Conference proceedings
Menzo 2008	No relevant outcomes reported
Meshher 2013	No results for women with relevant HPV and cytology status
Michelli 2011	No results for women with relevant HPV and cytology status
Michelli 2013	No relevant outcomes reported
Mijit 2015	No results for women with relevant HPV and cytology status
Min 2015	Conference proceedings
Moarcas 2014	No results for women with relevant HPV and cytology status
Monsonogo 2008	No results for women with relevant HPV and cytology status

Monsonogo 2008b	No results for women with relevant HPV and cytology status
Monsonogo 2011	No results for women with relevant HPV and cytology status
Monsonogo 2012	No results for women with relevant HPV and cytology status
Moore 2010	No results for women with relevant HPV and cytology status
Moss 2006	No results for women with relevant HPV and cytology status
Moy 2010	No results for women with relevant HPV and cytology status
Musa 2011	No relevant outcomes reported
Muwonge 2014	No results for women with relevant HPV and cytology status
Nakamura 2015	Colposcopy referral population and unclear if treatment follow-up patients excluded
Nam 2008	Participants women with CIN diagnosis
Namugenyi 2013	No results for women with relevant HPV and cytology status
Nasioutziki 2011	No results for women with relevant HPV and cytology status
Naucler 2007	No results for women with relevant HPV and cytology status
Naucler 2007 Human	No results for women with relevant HPV and cytology status
Naucler 2009	No results for women with relevant HPV and cytology status
Ngugi 2015	No results for women with relevant HPV and cytology status
Niccolai 2013	No results for women with relevant HPV and cytology status
Nomelini 2007	No results for women with relevant HPV and cytology status
Nomelini 2012	No results for women with relevant HPV and cytology status
Nygard 2014	No results for women with relevant HPV and cytology status
O'Connor 2008	No results for women with relevant HPV and cytology status
Ogilvie 2010	No results for women with relevant HPV and cytology status
Ogilvie 2012	No results for women with relevant HPV and cytology status
Okadome 2014	Participants women with CIN2 histology
Ovestad 2011	No results for women with relevant HPV and cytology status
Paba 2012	No results for women with relevant HPV and cytology status
Pacchiarotti 2014	No results for women with relevant HPV and cytology status
Padalko 2013	No results for women with relevant HPV and cytology status
Paengchit 2014	No comparator
Pajtlar 2010	No results for women with relevant HPV and cytology status
Pan 2013	No results for women with relevant HPV and cytology status
Pan 2014	No results for women with relevant HPV and cytology status
Pasquale 2015	No results for women with relevant HPV and cytology status
Perksanusak 201	No results for women with relevant HPV and cytology status
Perrons 2005	No results for women with relevant HPV and cytology status
Petry 2011	No results for women with relevant HPV and cytology status
Petry 2012	No results for women with relevant HPV and cytology status
Petry 2013	No results for women with relevant HPV and cytology status
Piccolo 2008	No results for women with relevant HPV and cytology status
Pity 2012	No results for women with relevant HPV and cytology status
Possati-Resende 2015	No results for women with relevant HPV and cytology status
Pouliakis 2015	No results for women with relevant HPV and cytology status
Pretorius 2013	No results for women with relevant HPV and cytology status
Pruski 2012	No results for women with relevant HPV and cytology status



Qiao 2014	No results for women with relevant HPV and cytology status
Rabelo-Santos 2009	No results for women with relevant HPV and cytology status
Ramzan 2015	Review
Rao 2013	No results for women with relevant HPV and cytology status
Ratnam 2010	No results for women with relevant HPV and cytology status
Ratnam 2011	No results for women with relevant HPV and cytology status
Rebolj 2011	Commentary
Rebolj 2014	No results for women with relevant HPV and cytology status
Rebolj 2015	No results for women with relevant HPV and cytology status
Reid 2015	No results for women with relevant HPV and cytology status
Riera 2013	Conference proceedings
Rijkaart 2012	No results for women with relevant HPV and cytology status
Rijkaart 2012	No results for women with relevant HPV and cytology status
Rijkaart 2012	No relevant comparator similarly followed up
Rodriguez 2008	No results for women with relevant HPV and cytology status
Rodriguez 2010	No results for women with relevant HPV and cytology status
Rokita 2012	No results for women with relevant HPV and cytology status
Ronco 2010	No results for women with relevant HPV and cytology status
Ronco 2014	No results for women with relevant HPV and cytology status
Saccardi 2014	No results for women with relevant HPV and cytology status
Safaeian 2009	No results for women with relevant HPV and cytology status
Sahasrabuddhe 2014	No results for women with relevant HPV and cytology status
Sahasrabuddhe 2014b	No results for women with relevant HPV and cytology status
Samarawardana 2010	No results for women with relevant HPV and cytology status
Sankaranarayanan 2009	No results for women with relevant HPV and cytology status
Saslow 2012	No results for women with relevant HPV and cytology status
Sauer 2011	Conference proceedings
Sauter 2014	No results for women with relevant HPV and cytology status
Schettino 2014	No results for women with relevant HPV and cytology status
Schiffman 2011	Discussion article
Schiffman 2011b	No results for women with relevant HPV and cytology status
Schiffman 2015	No relevant comparators similarly followed up
Schiffman 2015a	No results for women with relevant HPV and cytology status
Schiffman 2015b	No results for women with relevant HPV and cytology status
Schlichte 2014	Review
Schnatz 2009	Not relevant population - AGC cytology
Sellors 2011	No results for women with relevant HPV and cytology status
Selvaggi 2015	No results for women with relevant HPV and cytology status
Seo 2006	No results for women with relevant HPV and cytology status
Shahid 2015	No results for women with relevant HPV and cytology status
Shastri 2005	No results for women with relevant HPV and cytology status
Shen 2013	No results for women with relevant HPV and cytology status
Shen-Gunther 2011	Does not discuss multiple infections
Shi 2009	No results for women with relevant HPV and cytology status

Shipitsyna 2011	No results for women with relevant HPV and cytology status
Shwe 2015	No results for women with relevant HPV and cytology status
Siebers 2014	No results for women with relevant HPV and cytology status
Sigurdsson 2009	No results for women with relevant HPV and cytology status
Sikon 2014	Conference proceedings
Silva 2014	No relevant outcomes reported
Sodhani 2006	Not a screening population - women with non specific gynecological symptoms
Sole-Sedeno 2015	Conference proceedings
Solomon 2015	Conference proceedings
Song 2006	No results for women with relevant HPV and cytology status
Sorboye 2011	No results for women with relevant HPV and cytology status
Sorbye 2013	No results for women with relevant HPV and cytology status
Sorbye 2014	No results for women with relevant HPV and cytology status
Soto 2014	No colposcopy/biopsy/histology results
Spathis 2012	No results for women with relevant HPV and cytology status
Spinillo 2009	No results for women with relevant HPV and cytology status
Spinillo 2014	No results for women with relevant HPV and cytology status
Spinillo 2014b	No results for women with relevant HPV and cytology status
Stanculescu 2013	No results for women with relevant HPV and cytology status
Stanczuk 2015	No results for women with relevant HPV and cytology status
Stoler 2011	No results for women with relevant HPV and cytology status
Stoler 2013	No results for women with relevant HPV and cytology status
Stoler 2015	Commentary
Sultana 2015	No outcomes of interest
Supho 2014	No comparator
Sycuro 2008	No results for women with relevant HPV and cytology status
Szarewski 2007	No results for women with relevant HPV and cytology status
Szarewski 2008	No results for women with relevant HPV and cytology status
Szarewski 2011	Commentary
Szarewski 2012	No results for women with relevant HPV and cytology status
Tempfer 2007	No results for women with relevant HPV and cytology status
Tinelli 2013	Participants either women at high-risk of HPV infection or women who have been treated for CIN2+ diagnosis
Tokmak 2014	No results for women with relevant HPV and cytology status
Tornello 2007	No results for women with relevant HPV and cytology status
Tornello 2011	No relevant outcomes reported
Trimble 2015	Population women with CIN2/3 diagnosis
Trope 2012	No results for women with relevant HPV and cytology status
Tsai 2009	No results for women with relevant HPV and cytology status
Tsiodras 2010	No results for women with relevant HPV and cytology status
Tsiodras 2011	Conference proceedings
Tsompou 2011	Results for only a subgroup of women with relevant HPV and cytology status
Twiggs 2011	Discussion article
Twu 2011	No results for women with relevant HPV and cytology status



Uijterwaal 2014	No results for women with relevant HPV and cytology status
Uijterwaal 2015	No relevant comparator
Ursu 2011	No relevant outcomes reported
van der Marel 2014	No results for women with relevant HPV and cytology status
van der Marel 2015	No results for women with relevant HPV and cytology status
Van Le 2015	Summary of a published study
Verdoodt 2013	No results for women with relevant HPV and cytology status
Vidal 2013	Conference proceedings
Vidal 2014	No results for women with relevant HPV and cytology status
Vink 2014	Modelling effect of increasing screening interval
Wahlstrom 2007	No results for women with relevant HPV and cytology status
Walavalkar 2015	No results for women with relevant HPV and cytology status
Walker 2006	Participants women with cytological abnormalities and histology < CIN2
Walmer 2013	No results for women with relevant HPV and cytology status
Wang 2009	No results for women with relevant HPV and cytology status
Wang 2013	No results for women with relevant HPV and cytology status
Watson 2015	No results for women with relevant HPV and cytology status
Wei 2014	No results for women with relevant HPV and cytology status
Wentzensen 2009	Analysis restricted to women with CIN
Wentzensen 2009b	No results for women with relevant HPV and cytology status
Wentzensen 2010	No results for women with relevant HPV and cytology status
Wentzensen 2015	No results for women with relevant HPV and cytology status
Werner 2007	No results for women with relevant HPV and cytology status
Wheeler 2006	No results for women with relevant HPV and cytology status
Wheeler 2009	No relevant outcomes reported
Wheeler 2014	No results for women with relevant HPV and cytology status
White 2013	No results for women with relevant HPV and cytology status
Whitlock 2011	No results for women with relevant HPV and cytology status
Winer 2005	No results for women with relevant HPV and cytology status
Winsley 2014	No results for women with relevant HPV and cytology status
Wong 2008	No relevant outcomes reported
Woodard 2014	Conference proceedings
Woodard 2015	No results for women with relevant HPV and cytology status
Wright 2006	No results for women with relevant HPV and cytology status
Wright 2007a	Consensus guidelines (superseded)
Wright 2007b	Consensus guidelines (superseded)
Wright 2011	No longitudinal follow-up
Wright 2013	Conference proceedings
Wright 2014	Conference proceedings
Wright 2014	No results for women with relevant HPV and cytology status
Wright 2015	No comparable comparative data
Wu 2005	No results for women with relevant HPV and cytology status
Wu 2010	No results for women with relevant HPV and cytology status
Wu 2012	No results for women with relevant HPV and cytology status

Yao 2014	Conference proceedings
Ye 2010	No results for women with relevant HPV and cytology status
Yetimalar 2011	No results for women with relevant HPV and cytology status
You 2013	No results for women with relevant HPV and cytology status
Yuan 2011	No results for women with relevant HPV and cytology status
Zappacosta 2013	No results for women with relevant HPV and cytology status
Zeferino 2011	No results for women with relevant HPV and cytology status
Zeng 2011	No results for women with relevant HPV and cytology status
Zhang 2013	No results for women with relevant HPV and cytology status
Zhang 2014	Participants women with persistent HPV infection
Zhao 2013	No results for women with relevant HPV and cytology status
Zorzi 2013	No results for women with relevant HPV and cytology status

### References: Excluded Studies

1. Adamopoulou M, Kalkani E, Charvalos E et al. Comparison of cytology, colposcopy, HPV typing and biomarker analysis in cervical neoplasia. *Anticancer Research*. 2009;29:3401-3409.
2. Agodi A, Barchitta M, La RN et al. Human papillomavirus infection: low-risk and high-risk genotypes in women in Catania, Sicily. *Int J Gynecol Cancer*. 2009;19:1094-1098.
3. Agorastos T, Dinas K, Lloveras B et al. Human papillomavirus testing for primary screening in women at low risk of developing cervical cancer. The Greek experience. *Gynecol Oncol*. 2005;96:714-720.
4. Agorastos T, Chatzistamatiou K, Katsamagkas T et al. Primary screening for cervical cancer based on high-risk human papillomavirus (HPV) detection and HPV 16 and HPV 18 genotyping, in comparison to cytology. *PLoS ONE*. 2015;10.
5. Aksu M, Altinta+f A, Nazik H et al. One year follow up results of one hundred ASCUS patients. *Int J Gynecol Cancer*. 2011;21:S310.
6. Alameda F, Hernandez E, Lloveras B et al. HPV testing in ΓÇËatypical cells cannot exclude high grade lesionΓÇÏ (ASC-H). *Cytopathology*. 2014;25:56.
7. Allia E, Ronco G, Coccia A et al. Interpretation of p16(INK4a) /Ki-67 dual immunostaining for the triage of human papillomavirus-positive women by experts and nonexperts in cervical cytology. *Cancer Cytopathol*. 2015;123:212-218.
8. Almonte M, Ferreccio C, Winkler JL et al. Cervical screening by visual inspection, HPV testing, liquid-based and conventional cytology in Amazonian Peru. *International journal of cancer*. 2007;121:796-802.
9. Annappa R, Goodfellow J, Chalhoub T et al. High risk human papillomavirus triage. *BJOG Int J Obstet Gynaecol*. 2015;122:131.
10. Antonishyn NA, Horsman GB, Kelln RA et al. The impact of the distribution of human papillomavirus types and associated high-risk lesions in a colposcopy population for monitoring vaccine efficacy. *Archives of Pathology & Laboratory Medicine*. 2008;132:54-60.
11. Antonishyn NA, Horsman GB, Kelln RA et al. Human papillomavirus typing and viral gene expression analysis for the triage of women with abnormal results from papanicolaou test smears to colposcopy. *Archives of Pathology & Laboratory Medicine*. 2009;133:1577-1586.

12. Arbyn M, Buntinx F, Van RM et al. Virologic versus cytologic triage of women with equivocal Pap smears: a meta-analysis of the accuracy to detect high-grade intraepithelial neoplasia. *Journal of the National Cancer Institute*. 2004;96:280-293.
13. Arbyn M, Martin-Hirsch P, Buntinx F et al. Triage of women with equivocal or low-grade cervical cytology results: a meta-analysis of the HPV test positivity rate. *Journal of Cellular & Molecular Medicine*. 2009;13:648-659.
14. Arbyn M, Ronco G, Anttila A et al. Evidence regarding human papillomavirus testing in secondary prevention of cervical cancer. [Review][Erratum appears in *Vaccine*. 2013 Dec 16;31(52):6266]. *Vaccine*. 2012;30S:F88-F99.
15. Atkins KA, Jeronimo J, Stoler MH et al. Description of patients with squamous cell carcinoma in the atypical squamous cells of undetermined significance/low-grade squamous intraepithelial lesion triage study. *Cancer*. 2006;108:212-221.
16. Ault KA, Joura EA, Kjaer SK et al. Adenocarcinoma in situ and associated human papillomavirus type distribution observed in two clinical trials of a quadrivalent human papillomavirus vaccine. *International journal of cancer*. 2011;128:1344-1353.
17. Badano I, Pedrozo RW, Ruiz Diaz LS et al. Human papillomavirus (HPV) detection and Papanicolaou cytology in low-resource women in Posadas city, Misiones, Argentina. *Revista Argentina de Microbiologia*. 2011;43:263-267.
18. Bais AG, Rebolj M, Snijders PJ et al. Triage using HPV-testing in persistent borderline and mildly dyskaryotic smears: proposal for new guidelines. *International journal of cancer*. 2005;116:122-129.
19. Balbi G, Napolitano A, Giordano F et al. Role of the association of high-risk HPV identified by real-time PCR in cervical preneoplastic lesions. *European Journal of Gynaecological Oncology*. 2012;33:467-471.
20. Bapir M, Arutchelvam S, Doxford L et al. Review of outcome of patients referred to colposcopy with low grade smears after introduction of HPV triage. *BJOG Int J Obstet Gynaecol*. 2013;120:54.
21. Barcelos AC, Michelin MA, Adad SJ et al. Atypical squamous cells of undetermined significance: Bethesda classification and association with Human Papillomavirus. *Infectious Diseases in Obstetrics & Gynecology*. 2011;2011:904674.
22. Barzon L, Militello V, Lavezzo E et al. Human papillomavirus genotyping by 454 next generation sequencing technology. *J Clin Virol*. 2011;52:93-97.
23. Basu P, Chandna P, Bamezai RN et al. MassARRAY spectrometry is more sensitive than PreTect HPV-Proofer and consensus PCR for type-specific detection of high-risk oncogenic human papillomavirus genotypes in cervical cancer. *J Clin Microbiol*. 2011;49:3537-3544.
24. Bekkers RL, Hanselaar AG, Melchers WJ et al. Management of patients with two consecutive ASC-US smears. *European Journal of Gynaecological Oncology*. 2004;25:677-681.
25. Belinson JL, Wu R, Belinson SE et al. A population-based clinical trial comparing endocervical high-risk HPV testing using hybrid capture 2 and Cervista from the SHENCCAST II Study. *American Journal of Clinical Pathology*. 2011;135:790-795.
26. Belinson SE, Wulan N, Li R et al. SNIPER: a novel assay for human papillomavirus testing among women in Guizhou, China. *Int J Gynecol Cancer*. 2010;20:1006-1010.
27. Bello BD, Spinillo A, Alberizzi P et al. Cervical infections by multiple human papillomavirus (HPV) genotypes: Prevalence and impact on the risk of precancerous epithelial lesions. *Journal of Medical Virology*. 2009;81:703-712.

28. Benevolo M, Terrenato I, Mottolese M et al. Diagnostic and prognostic validity of the human papillomavirus E6/E7 mRNA test in cervical cytological samples of HC2-positive patients. *Cancer Causes Control*. 2011;22:869-875.
29. Benevolo M, Vocaturo A, Caraceni D et al. Sensitivity, specificity, and clinical value of human papillomavirus (HPV) E6/E7 mRNA assay as a triage test for cervical cytology and HPV DNA test. *J Clin Microbiol*. 2011;49:2643-2650.
30. Benoy IH, Vanden Broeck D, Ruymbeke MJ et al. Prior knowledge of HPV status improves detection of CIN2+ by cytology screening. *American Journal of Obstetrics & Gynecology*. 2011;205:569-7.
31. Bergeron C, Giorgi-Rossi P, Cas F et al. Informed cytology for triaging HPV-positive women: substudy nested in the NTCC randomized controlled trial. *J Natl Cancer Inst*. 2015;107.
32. Berkhof J, de Bruijne MC, Zielinski GD et al. Evaluation of cervical screening strategies with adjunct high-risk human papillomavirus testing for women with borderline or mild dyskaryosis. *International journal of cancer*. 2006;118:1759-1768.
33. Bhatla N, Mukhopadhyay A, Kriplani A et al. Evaluation of adjunctive tests for cervical cancer screening in low resource settings. *Indian Journal of Cancer*. 2007;44:51-55.
34. Bhatla N, Dar L, Rajkumar PA et al. Human papillomavirus-type distribution in women with and without cervical neoplasia in north India. *International Journal of Gynecological Pathology*. 2008;27:426-430.
35. Bian M-L, Cheng J-Y, Li M et al. Evaluation of the detection of 14 high-risk human papillomaviruses with HPV 16 and HPV 18 genotyping for cervical cancer screening. *Exp Ther Med*. 2013;6:1332-1336.
36. Bjerre P, Silfverdal L, Dillner L et al. A randomized trial of basing treatment on human papillomavirus and/or cytology results in low-grade cervical lesion triage. *American Journal of Obstetrics & Gynecology*. 2008;199:24-27.
37. Blatt AJ, Kennedy R, Luff RD et al. Comparison of cervical cancer screening results among 256,648 women in multiple clinical practices. *Cancer Cytopathol*. 2015;123:282-288.
38. Boardman LA, Weitzen S, Stanko C. Atypical squamous cells of undetermined significance, human papillomavirus, and cervical intraepithelial neoplasia 2 or 3 in adolescents: ASC-US, age, and high-grade cervical neoplasia. *J Lower Genital Tract Dis*. 2006;10:140-145.
39. Boldrini NT, Freitas LB, Coutinho AR et al. High-grade cervical lesions among women attending a reference clinic in Brazil: associated factors and comparison among screening methods. *PLoS ONE [Electronic Resource]*. 2014;9:e102169.
40. Bollmann M, Varnai AD, Griefingholt H et al. Predicting treatment outcome in cervical diseases using liquid-based cytology, dynamic HPV genotyping and DNA cytometry. *Anticancer Research*. 2006;26:1439-1446.
41. Bourgain C, Bagar W, Willocx F. A 10-year follow-up of ASC-US and ASC-H cytological diagnosis with Surepath-«. *Cytopathology*. 2011;22:61.
42. Bratti MC, Rodriguez AC, Schiffman M et al. Description of a seven-year prospective study of human papillomavirus infection and cervical neoplasia among 10000 women in Guanacaste, Costa Rica. *Pan American Journal of Public Health*. 2004;15:75-89.
43. Brink AA, Meijer CJ, Wiegerinck MA et al. High concordance of results of testing for human papillomavirus in cervicovaginal samples collected by two methods, with comparison of a novel self-sampling device to a conventional endocervical brush. *J Clin Microbiol*. 2006;44:2518-2523.

44. Bulk S, Bulkmans NW, Berkhof J et al. Risk of high-grade cervical intra-epithelial neoplasia based on cytology and high-risk HPV testing at baseline and at 6-months.[Erratum appears in *Int J Cancer*. 2007 Oct 15;121(8):1873]. *International journal of cancer*. 2007;121:361-367.
45. Bulkmans NW, Berkhof J, Rozendaal L et al. Human papillomavirus DNA testing for the detection of cervical intraepithelial neoplasia grade 3 and cancer: 5-year follow-up of a randomised controlled implementation trial. *Lancet*. 2007;370:1764-1772.
46. Bulkmans NW, Berkhof J, Bulk S et al. High-risk HPV type-specific clearance rates in cervical screening. *British journal of cancer*. 2007;96:1419-1424.
47. Campari C, Fedato C, Petrelli A et al. HPV prevalence and risk of pre-cancer and cancer in regular immigrants in Italy: results from HPV DNA test-based screening pilot programs. *Infectious Agents & Cancer [Electronic Resource]*. 2015;10:14.
48. Canda MT, Demir N, Sezer O et al. Clinical results of the liquid-based cervical cytology tool, Liqueu-PREP, in comparison with conventional smears for detection of squamous cell abnormalities. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2009;10:399-402.
49. Carozzi F, Cecchini S, Confortini M et al. Role of P16(INK4a) expression in identifying CIN2 or more severe lesions among HPV-positive patients referred for colposcopy after abnormal cytology. *Cancer*. 2006;108:119-123.
50. Carozzi F, Visioli CB, Confortini M et al. hr-HPV testing in the follow-up of women with cytological abnormalities and negative colposcopy. *British journal of cancer*. 2013;109:1766-1774.
51. Carozzi F, Gillio-Tos A, Confortini M et al. Risk of high-grade cervical intraepithelial neoplasia during follow-up in HPV-positive women according to baseline p16-INK4A results: a prospective analysis of a nested substudy of the NTCC randomised controlled trial. *Lancet Oncology*. 2013;14:168-176.
52. Carozzi FM, Confortini M, Cecchini S et al. Triage with human papillomavirus testing of women with cytologic abnormalities prompting referral for colposcopy assessment. *Cancer*. 2005;105:2-7.
53. Carter JS, Sage YH, Vragovic O et al. Effect of patient age on outcomes and compliance in women with minimally abnormal pap tests. *Journal of Reproductive Medicine*. 2010;55:351-356.
54. Castle PE, Schiffman M. Women with a positive HPV test can safely have yearly surveillance rather than immediate colposcopy. *Evid -based Obstet Gynecol*. 2004;6:150-151.
55. Castle PE, Stoler MH, Solomon D et al. The relationship of community biopsy-diagnosed cervical intraepithelial neoplasia grade 2 to the quality control pathology-reviewed diagnoses: an ALTS report. *American Journal of Clinical Pathology*. 2007;127:805-815.
56. Castle PE, Cox JT, Schiffman M et al. Factors influencing histologic confirmation of high-grade squamous intraepithelial lesion cytology. *Obstetrics & Gynecology*. 2008;112:637-645.
57. Castle PE, Rodriguez AC, Burk RD et al. Neither one-time negative screening tests nor negative colposcopy provides absolute reassurance against cervical cancer. *International journal of cancer*. 2009;125:1649-1656.
58. Castle PE, Wentzensen N, Wheeler CM et al. Effect of the number of biopsies on the subsequent acquisition of new human papillomavirus infections. *Obstetrics & Gynecology*. 2009;114:1057-1062.
59. Castle PE, Rodriguez AC, Burk RD et al. Short term persistence of human papillomavirus and risk of cervical precancer and cancer: population based cohort study. *BMJ*. 2009;339:b2569.

60. Castle PE, Schiffman M, Wheeler CM et al. Evidence for frequent regression of cervical intraepithelial neoplasia-grade 2. *Obstetrics & Gynecology*. 2009;113:18-25.
61. Castle PE, Fetterman B, Thomas CJ et al. The age-specific relationships of abnormal cytology and human papillomavirus DNA results to the risk of cervical precancer and cancer.[Erratum appears in *Obstet Gynecol*. 2010 Sep;116(3):775-7]. *Obstetrics & Gynecology*. 2010;116:76-84.
62. Castle PE, Stoler MH, Wright TC, Jr. et al. Performance of carcinogenic human papillomavirus (HPV) testing and HPV16 or HPV18 genotyping for cervical cancer screening of women aged 25 years and older: a subanalysis of the ATHENA study. *Lancet Oncology*. 2011;12:880-890.
63. Castle PE, Gutierrez EC, Leitch SV et al. Evaluation of a new DNA test for detection of carcinogenic human papillomavirus. *J Clin Microbiol*. 2011;49:3029-3032.
64. Castle PE, Gage JC, Wheeler CM et al. The clinical meaning of a cervical intraepithelial neoplasia grade 1 biopsy. *Obstetrics & Gynecology*. 2011;118:1222-1229.
65. Castle PE, Glass AG, Rush BB et al. Clinical human papillomavirus detection forecasts cervical cancer risk in women over 18 years of follow-up. *J Clin Oncol*. 2012;30:3044-3050.
66. Castle PE, Eaton B, Reid J et al. Comparison of human papillomavirus detection by Aptima HPV and cobas HPV tests in a population of women referred for colposcopy following detection of atypical squamous cells of undetermined significance by Pap cytology. *J Clin Microbiol*. 2015;53:1277-1281.
67. Castle PE, Reid J, Dockter J et al. The reliability of high-risk human papillomavirus detection by Aptima HPV assay in women with ASC-US cytology. *J Clin Virol*. 2015;69:52-55.
68. Castle PE, Smith KM, Davis TE et al. Reliability of the Xpert HPV assay to detect high-risk human papillomavirus DNA in a colposcopy referral population. *American Journal of Clinical Pathology*. 2015;143:126-133.
69. Cattani P, Zannoni GF, Ricci C et al. Clinical performance of human papillomavirus E6 and E7 mRNA testing for high-grade lesions of the cervix. *J Clin Microbiol*. 2009;47:3895-3901.
70. Chao A, Hsu KH, Lai CH et al. Cervical cancer screening program integrating Pap smear and HPV DNA testing: a population-based study. *International journal of cancer*. 2008;122:2835-2841.
71. Chao A, Chang CJ, Lai CH et al. Incidence and outcome of acquisition of human papillomavirus infection in women with normal cytology--a population-based cohort study from Taiwan. *International journal of cancer*. 2010;126:191-198.
72. Chen CA, Liu CY, Chou HH et al. The distribution and differential risks of human papillomavirus genotypes in cervical preinvasive lesions: A Taiwan Cooperative Oncologic Group Study. *Int J Gynecol Cancer*. 2006;16:1801-1808.
73. Chen X, Wallin KL, Duan M et al. Prevalence and genotype distribution of cervical human papillomavirus (HPV) among women in urban Tianjin, China. *Journal of Medical Virology*. 2015;87:1966-1972.
74. Chevarie-Davis M, Ramanakumar AV, Ferenczy A et al. Assessment of the performance of algorithms for cervical cancer screening: evidence from the Ludwig-McGill cohort study. *Gynecol Oncol*. 2013;128:415-419.
75. Chiang YC, Cheng WF, Chen YL et al. High-risk human papillomavirus, other than type 16/18, in predominantly older Taiwanese women with high-grade cervical preinvasive lesions. *Taiwanese Journal of Obstetrics & Gynecology*. 2013;52:222-226.

76. Chivukula M, Shidham VB. ASC-H in Pap test- definitive categorization of cytomorphological spectrum. *CytoJournal*. 2006;3.
77. Cho HW, So KA, Lee JK et al. Type-specific persistence or regression of human papillomavirus genotypes in women with cervical intraepithelial neoplasia 1: A prospective cohort study. *Obstetrics & Gynecology Science*. 2015;58:40-45.
78. Chranioti A, Spathis A, Aga E et al. Comparison of two commercially available methods for HPV genotyping: CLART HPV2 and Linear Array HPV Genotyping tests. *Analytical & Quantitative Cytology & Histology*. 2012;34:257-263.
79. Cilingir IU, Bengisu E, Agacfidan A et al. Microarray detection of human papilloma virus genotypes among Turkish women with abnormal cytology at a colposcopy unit. *Journal of the Turkishgerman Gynecological Association*. 2013;14:23-27.
80. Ciotti M, Sesti F, Paba P et al. Human papillomavirus (HPV) testing in the management of women with abnormal Pap smears. Experience of a colposcopy referral clinic. *European Journal of Gynaecological Oncology*. 2004;25:577-584.
81. Clad A, Reuschenbach M, Weinschenk J et al. Performance of the Aptima high-risk human papillomavirus mRNA assay in a referral population in comparison with Hybrid Capture 2 and cytology. *J Clin Microbiol*. 2011;49:1071-1076.
82. Clarke M, Boyle S, Burk R et al. Performance of cytology and HPV16/18 genotyping among a large cohort of HPV positive women aged 30 years and older. *Sex Transm Dis*. 2014;41:S76-S77.
83. Cobo F, Concha A, Ortiz M. Human Papillomavirus (HPV) Type Distribution in Females with Abnormal Cervical Cytology. A Correlation with Histological Study. *The Open Virology Journal [electronic resource]*. 2009;3:60-66.
84. Confortini M, Carozzi F, Zappa M et al. Human papillomavirus infection and risk factors in a cohort of Tuscan women aged 18-24: results at recruitment. *BMC Infectious Diseases*. 2010;10:157.
85. Corneanu LM, Stanculescu D, Corneanu C. HPV and cervical squamous intraepithelial lesions: clinicopathological study. *Romanian Journal of Morphology & Embryology*. 2011;52:89-94.
86. Correa FM, Russomano FB, Oliveira CA. Colposcopic triage methods for detecting cervical intraepithelial neoplasia grade 3 after cytopathological diagnosis of low-grade squamous intraepithelial lesion: a systematic review on diagnostic tests. [Review]. *Sao Paulo Medical Journal = Revista Paulista de Medicina*. 2012;130:44-52.
87. Cottier O, Sahli R, Mihaescu A et al. Clinical follow-up of women infected with human papillomavirus-16, either alone or with other human papillomavirus types: identification of different risk groups. *American Journal of Obstetrics & Gynecology*. 2009;200:286.
88. Cotton S, Sharp L, Little J et al. The role of human papillomavirus testing in the management of women with low-grade abnormalities: multicentre randomised controlled trial. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2010;117:645-659.
89. Cox JT, Castle PE, Behrens CM et al. Comparison of cervical cancer screening strategies incorporating different combinations of cytology, HPV testing, and genotyping for HPV 16/18: results from the ATHENA HPV study. *American Journal of Obstetrics & Gynecology*. 2013;208:184.
90. Cruickshank ME, Cotton SC, Sharp L et al. Management of women with low grade cytology: how reassuring is a normal colposcopy examination? *BJOG: An International Journal of Obstetrics & Gynaecology*. 2015;122:380-386.

91. Crum CP, Beach KJ, Hedley ML et al. Dynamics of human papillomavirus infection between biopsy and excision of cervical intraepithelial neoplasia: results from the ZYC101a protocol. *Journal of infectious diseases*. 2004;189:1348-1354.
92. Cuschieri K, Cubie H, Graham C et al. Clinical performance of RNA and DNA based HPV testing in a colposcopy setting: Influence of assay target, cut off and age. *J Clin Virol*. 2014;59:104-108.
93. Cuzick J, Arbyn M, Sankaranarayanan R et al. Overview of human papillomavirus-based and other novel options for cervical cancer screening in developed and developing countries. [Review] [91 refs]. *Vaccine*. 2008;26S:K29-K41.
94. Cuzick J, Ambroisine L, Cadman L et al. Performance of the Abbott RealTime high-risk HPV test in women with abnormal cervical cytology smears. *Journal of Medical Virology*. 2010;82:1186-1191.
95. Cuzick J, Thomas CJ, Zhang G et al. Human papillomavirus testing for triage of women with low-grade squamous intraepithelial lesions. *International journal of cancer*. 2013;132:959-966.
96. Dalla PP, Giorgi RP, Collina G et al. The risk of false-positive histology according to the reason for colposcopy referral in cervical cancer screening: a blind revision of all histologic lesions found in the NTCC trial. *American Journal of Clinical Pathology*. 2008;129:75-80.
97. Dalstein V, Riethmuller D, Sautiere JL et al. Detection of cervical precancer and cancer in a hospital population; benefits of testing for human papillomavirus.[Erratum appears in Eur J Cancer. 2005 Jan;41(2):334]. *European Journal of Cancer*. 2004;40:1225-1232.
98. Dane C, Batmaz G, Dane B et al. Screening properties of human papillomavirus testing for predicting cervical intraepithelial neoplasia in atypical squamous cells of undetermined significance and low-grade squamous intraepithelial lesion smears: a prospective study. *Annals of Diagnostic Pathology*. 2009;13:73-77.
99. Daponte A, Pournaras S, Mademtzis I et al. Evaluation of high-risk human papillomavirus types PCR detection in paired urine and cervical samples of women with abnormal cytology. *J Clin Virol*. 2006;36:189-193.
100. De VH, Claeys P, Njiru S et al. Comparison of pap smear, visual inspection with acetic acid, human papillomavirus DNA-PCR testing and cervicography. *International Journal of Gynaecology & Obstetrics*. 2005;89:120-126.
101. Del MA, Frayle-Salamanca H, Trevisan R et al. Triage of women with atypical squamous cells of undetermined significance (ASC-US): results of an Italian multicentric study. *Gynecol Oncol*. 2010;117:77-81.
102. Del MA, Frayle H, Ferro A et al. Cervical cancer screening by high risk HPV testing in routine practice: results at one year recall of high risk HPV-positive and cytology-negative women. *Journal of Medical Screening*. 2014;21:30-37.
103. Denny L, Kuhn L, Hu CC et al. Human papillomavirus-based cervical cancer prevention: long-term results of a randomized screening trial. *Journal of the National Cancer Institute*. 2010;102:1557-1567.
104. Depuydt CE, Makar AP, Ruymbeke MJ et al. BD-ProExC as adjunct molecular marker for improved detection of CIN2+ after HPV primary screening. *Cancer Epidemiology, Biomarkers & Prevention*. 2011;20:628-637.
105. Desteli GA, Demiralay E, Gursu T et al. Detection of HPV positivity by immunohistochemistry in colposcopic cervical biopsies with a cytological diagnosis of ASCUS. *Turk Patoloji Dergisi*. 2014;30:166-170.



106. Diaz-Montes TP, Farinola MA, Zahurak ML et al. Clinical utility of atypical glandular cells (AGC) classification: cytohistologic comparison and relationship to HPV results. *Gynecol Oncol.* 2007;104:366-371.
107. Dijkstra MG, van ND, Rijkaart DC et al. Primary hrHPV DNA testing in cervical cancer screening: how to manage screen-positive women? A POBASCAM trial substudy. *Cancer Epidemiology, Biomarkers & Prevention.* 2014;23:55-63.
108. Dillner J, Rebolj M, Birembaut P et al. Long term predictive values of cytology and human papillomavirus testing in cervical cancer screening: joint European cohort study. *BMJ.* 2008;337:a1754.
109. Dillner L, Kemetli L, Elfgrén K et al. Randomized healthservices study of human papillomavirus-based management of low-grade cytological abnormalities. *International journal of cancer.* 2011;129:151-159.
110. Dinc B, Rota S, Onan A et al. Prevalence of human papillomavirus (HPV) and HPV-16 genotyping by real-time PCR in patients with several cervical pathologies. *Brazilian Journal of Infectious Diseases.* 2010;14:19-23.
111. Ding X, Liu Z, Su J et al. Human papillomavirus type-specific prevalence in women referred for colposcopic examination in Beijing. *Journal of Medical Virology.* 2014;86:1937-1943.
112. Ding Z, Jiang C, Shore T et al. Outcome of cervical intraepithelial neoplasia 2 diagnosed by punch biopsy in 131 women. *Journal of Obstetrics & Gynaecology Research.* 2011;37:754-761.
113. Discacciati MG, da Silva ID, Villa LL et al. Prognostic value of DNA and mRNA e6/e7 of human papillomavirus in the evolution of cervical intraepithelial neoplasia grade 2. *Biomark Insights.* 2014;9:15-22.
114. Dockter J, Schroder A, Hill C et al. Clinical performance of the APTIMA HPV Assay for the detection of high-risk HPV and high-grade cervical lesions. *J Clin Virol.* 2009;45:Suppl-61.
115. Du H, Yi J, Wu R et al. A new PCR-based mass spectrometry system for high-risk HPV, part II: clinical trial. *American Journal of Clinical Pathology.* 2011;136:920-923.
116. Dufresne S, Sauthier P, Mayrand MH et al. Human papillomavirus (HPV) DNA triage of women with atypical squamous cells of undetermined significance with Amplicor HPV and Hybrid Capture 2 assays for detection of high-grade lesions of the uterine cervix. *J Clin Microbiol.* 2011;49:48-53.
117. Einstein MH, Martens MG, Garcia FA et al. Clinical validation of the Cervista HPV HR and 16/18 genotyping tests for use in women with ASC-US cytology. *Gynecol Oncol.* 2010;118:116-122.
118. Einstein MH, Garcia FA, Mitchell AL et al. Age-stratified performance of the Cervista HPV 16/18 genotyping test in women with ASC-US cytology. *Cancer Epidemiology, Biomarkers & Prevention.* 2011;20:1185-1189.
119. Elfgrén K, Rylander E, Radberg T et al. Colposcopic and histopathologic evaluation of women participating in population-based screening for human papillomavirus deoxyribonucleic acid persistence. *American journal of obstetrics and gynecology.* 2005;193:650-657.
120. Eltoum IA, Chhieng DC, Roberson J et al. Reflex human papilloma virus infection testing detects the same proportion of cervical intraepithelial neoplasia grade 2-3 in young versus elderly women. *Cancer.* 2005;105:194-198.

121. Filho LA, Utagawa ML, Shirata NK et al. Immunocytochemical expression of p16INK4A and Ki-67 in cytologically negative and equivocal pap smears positive for oncogenic human papillomavirus. *International Journal of Gynecological Pathology*. 2005;24:118-124.
122. Fokom-Domgue J, Combescure C, Fokom-Defo V et al. Performance of alternative strategies for primary cervical cancer screening in sub-Saharan Africa: systematic review and meta-analysis of diagnostic test accuracy studies. *BMJ*. 2015;351:h3084.
123. Froberg M, Johansson B, Hjerpe A et al. Human papillomavirus 'reflex' testing as a screening method in cases of minor cytological abnormalities. *British journal of cancer*. 2008;99:563-568.
124. Gage JC, Schiffman M, Solomon D et al. Comparison of measurements of human papillomavirus persistence for postcolposcopic surveillance for cervical precancerous lesions. *Cancer Epidemiology, Biomarkers & Prevention*. 2010;19:1668-1674.
125. Gage JC, Schiffman M, Solomon D et al. Risk of precancer determined by HPV genotype combinations in women with minor cytologic abnormalities. *Cancer Epidemiology, Biomarkers & Prevention*. 2013;22:1095-1101.
126. Gage JC, Katki HA, Schiffman M et al. The low risk of precancer after a screening result of human papillomavirus-negative/atypical squamous cells of undetermined significance papanicolaou and implications for clinical management. *Cancer Cytopathol*. 2014;122:842-850.
127. Geng L, You K, Guo Y et al. Study of 361 cases of cervical cytology interpreted as atypical squamous cells which cannot exclude high-grade squamous intraepithelial lesion (ASC-H). *Cytopathology*. 2011;22:48.
128. Geraets DT, van BR, Alonso I et al. Clinical evaluation of high-risk HPV detection on self-samples using the indicating FTA-elute solid-carrier cartridge. *J Clin Virol*. 2013;57:125-129.
129. Gilani SM, Mazzara PF. Cytohistologic correlation in premenopausal and postmenopausal women. *Acta Cytologica*. 2013;57:575-580.
130. Giorgi RP, Bisanzi S, Paganini I et al. Prevalence of HPV high and low risk types in cervical samples from the Italian general population: a population based study. *BMC Infectious Diseases*. 2010;10:214.
131. Giorgi RP, Chini F, Bisanzi S et al. Distribution of high and low risk HPV types by cytological status: a population based study from Italy. *Infectious Agents & Cancer [Electronic Resource]*. 2011;6:2.
132. Giorgi RP, Benevolo M, Vocaturo A et al. Prognostic value of HPV E6/E7 mRNA assay in women with negative colposcopy or CIN1 histology result: a follow-up study. *PLoS ONE [Electronic Resource]*. 2013;8:e57600.
133. Giovannelli L, Capra G, Lama A et al. Atypical squamous cells of undetermined significance-favour reactive compared to atypical squamous cells of undetermined significance-favour dysplasia: association with cervical intraepithelial lesions and human papillomavirus infection. *J Clin Virol*. 2005;33:281-286.
134. Girianelli VR, Thuler LC, Szklo M et al. Comparison of human papillomavirus DNA tests, liquid-based cytology and conventional cytology for the early detection of cervix uteri cancer. *European Journal of Cancer Prevention*. 2006;15:504-510.
135. Gold MA, Thomas MA, Huh WK et al. High-risk human papillomavirus detection in women with low-grade squamous intraepithelial lesions or higher-grade cytology using the Cervista HPV HR test. *J Lower Genital Tract Dis*. 2013;17:51-57.

136. Gonzalez-Bosquet E, Almagro MM, Mora I et al. Prevalence of human papilloma virus infection of the uterine cervix in women with abnormal cervical cytology. *European Journal of Gynaecological Oncology*. 2006;27:135-138.
137. Gonzalez-Bosquet E, Esteva C, Munoz-Almagro C et al. Identification of vaccine human papillomavirus genotypes in squamous intraepithelial lesions (CIN2-3). *Gynecol Oncol*. 2008;111:9-12.
138. Gonzalez-Losa MR, Rosado-Lopez I, Valdez-Gonzalez N et al. High prevalence of human papillomavirus type 58 in Mexican colposcopy patients. *J Clin Virol*. 2004;29:202-205.
139. Goodrich SK, Pretorius RG, Du H et al. Triage of women with negative cytology and positive high-risk HPV: an analysis of data from the SHENCCAST II/III studies. *J Lower Genital Tract Dis*. 2014;18:122-127.
140. Gravitt PE, Paul P, Katki HA et al. Effectiveness of VIA, Pap, and HPV DNA testing in a cervical cancer screening program in a peri-urban community in Andhra Pradesh, India. *PLoS ONE [Electronic Resource]*. 2010;5:e13711.
141. Grisaru D, Avidor B, Niv J et al. Pilot study of prevalence of high-risk human papillomavirus genotypes in Israeli Jewish women referred for colposcopic examination. *J Clin Microbiol*. 2008;46:1602-1605.
142. Guo Y, You K, Geng L et al. The clinical performance of APTIMA human papillomavirus and Hybrid Capture 2 assays in the triage of lesser abnormal cervical cytologies. *Journal of Gynecologic Oncology*. 2014;25:287-292.
143. Gurumurthy M, Cotton SC, Sharp L et al. Postcolposcopy management of women with histologically proven CIN 1: results from TOMBOLA. *J Lower Genital Tract Dis*. 2014;18:203-209.
144. Gutierrez-Delgado F, Marcial-Toledo S, Enriquez-Freire JM et al. Cervical cancer screening: Twelve years experience with high-risk population of southern Mexico. *J Clin Oncol*. 2014;32.
145. Haidopoulos D, Partsinevelos GA, Vlachos GD et al. p16 INK4A is a strong biomarker for cervical intraepithelial neoplasia and invasive cervical carcinoma: a reappraisal. *Reprod Sci*. 2009;16:685-693.
146. Haldorsen T, Skare GB, Ursin G et al. Results of delayed triage by HPV testing and cytology in the Norwegian Cervical Cancer Screening Programme. *Acta Oncologica*. 2015;54:200-209.
147. Halfon P, Benmoura D, Agostini A et al. Relevance of HPV mRNA detection in a population of ASCUS plus women using the NucliSENS EasyQ HPV assay. *J Clin Virol*. 2010;47:177-181.
148. Halfon P, Lindemann ML, Raimondo A et al. HPV genotype distribution according to severity of cervical neoplasia using the Digene HPV genotyping LQ test. *Archives of Virology*. 2013;158:1143-1149.
149. Hariri S, Markowitz LE, Bennett NM et al. Monitoring Effect of Human Papillomavirus Vaccines in US Population, Emerging Infections Program, 2008-2012. *Emerging Infectious Diseases*. 2015;21:1557-1561.
150. Hariri S, Bennett NM, Niccolai LM et al. Reduction in HPV 16/18-associated high grade cervical lesions following HPV vaccine introduction in the United States - 2008-2012. *Vaccine*. 2015;33:1608-1613.
151. HooKim K, Lieberman RW, Smola BM et al. "equivocal" high-risk HPV DNA tests performed on thinprep specimens after ASC-US diagnoses are associated with an increased incidence of CIN3: A cytologic/histologic review of 315 cases. *J Lower Genital Tract Dis*. 2014.

152. Hopenhayn C, Christian A, Christian WJ et al. Prevalence of human papillomavirus types in invasive cervical cancers from 7 US cancer registries before vaccine introduction. *J Lower Genital Tract Dis.* 2014;18:182-189.
153. Hou R, Xu C, Zhang S et al. Distribution of human papillomavirus genotype and cervical neoplasia among women with abnormal cytology in Beijing, China. *International Journal of Gynaecology & Obstetrics.* 2012;119:257-261.
154. Hoyer H, Scheungraber C, Kuehne-Heid R et al. Cumulative 5-year diagnoses of CIN2, CIN3 or cervical cancer after concurrent high-risk HPV and cytology testing in a primary screening setting. *International journal of cancer.* 2005;116:136-143.
155. Hu SY, Hong Y, Zhao FH et al. Prevalence of HPV Infection And Cervical Intraepithelial Neoplasia And Attitudes towards HPV Vaccination among Chinese Women Aged 18-25 in Jiangsu Province. *Chinese Journal of Cancer Research.* 2011;23:25-32.
156. Huang S, Erickson B, Tang N et al. Clinical performance of Abbott RealTime High Risk HPV test for detection of high-grade cervical intraepithelial neoplasia in women with abnormal cytology. *J Clin Virol.* 2009;45:Suppl-23.
157. Huang SL, Chao A, Hsueh S et al. Comparison between the Hybrid Capture II Test and an SPF1/GP6+ PCR-based assay for detection of human papillomavirus DNA in cervical swab samples. *J Clin Microbiol.* 2006;44:1733-1739.
158. Hughes AA, Glazner J, Barton P et al. A cost-effectiveness analysis of four management strategies in the determination and follow-up of atypical squamous cells of undetermined significance. *Diagnostic Cytopathology.* 2005;32:125-132.
159. Huh WK, Ault KA, Chelmow D et al. Use of primary high-risk human papillomavirus testing for cervical cancer screening: Interim clinical guidance. *J Lower Genital Tract Dis.* 2015;19:91-96.
160. Ibanez R, Autonell J, Sarda M et al. Protecting the underscreened women in developed countries: the value of HPV test. *BMC cancer.* 2014;14:574.
161. Iftner T, Becker S, Neis KJ et al. Head-to-Head Comparison of the RNA-Based Aptima Human Papillomavirus (HPV) Assay and the DNA-Based Hybrid Capture 2 HPV Test in a Routine Screening Population of Women Aged 30 to 60 Years in Germany. *J Clin Microbiol.* 2015;53:2509-2516.
162. Ikenberg H, Bergeron C, Schmidt D et al. Screening for cervical cancer precursors with p16/Ki-67 dual-stained cytology: results of the PALMS study. *Journal of the National Cancer Institute.* 2013;105:1550-1557.
163. Inoue M, Sakaguchi J, Sasagawa T et al. The evaluation of human papillomavirus DNA testing in primary screening for cervical lesions in a large Japanese population. *Int J Gynecol Cancer.* 2006;16:1007-1013.
164. Jakobsson M, Tarkkanen J, Auvinen E et al. Colposcopy referral rate can be reduced by high-risk human papillomavirus triage in the management of recurrent atypical squamous cells of undetermined significance or low-grade squamous intraepithelial lesion cytology in Finland. *International Journal of STD & AIDS.* 2012;23:485-489.
165. Jariene K, Vaitkiene D, Bartusevicius A et al. Prevalence of human papillomavirus types 16, 18, and 45 in women with cervical intraepithelial changes: associations with colposcopic and histological findings. *Medicina (Kaunas, Lithuania).* 2012;48:22-30.
166. Jentschke M, Soergel P, Hillemanns P. Evaluation of a multiplex real time PCR assay for the detection of human papillomavirus infections on self-collected cervicovaginal lavage samples. *Journal of Virological Methods.* 2013;193:131-134.

167. Jeronimo J, Bansil P, Valdez M et al. The Influence of Human Papillomavirus Genotypes on Visual Screening and Diagnosis of Cervical Precancer and Cancer. *J Lower Genital Tract Dis.* 2015;19:220-223.
168. Jiang L, Zeng Y, Li J et al. Performance of high-risk human papillomavirus testing in the triage of abnormal cervical cytology among Chinese younger women in Shanghai, China. *Asian Pacific Journal of Cancer Prevention: Apjcp.* 2011;12:2963-2967.
169. Johnson LR, Starkey CR, Palmer J et al. A comparison of two methods to determine the presence of high-risk HPV cervical infections. *American Journal of Clinical Pathology.* 2008;130:401-408.
170. Jones HW. Clinical validation of the cervista HPV HR and 16/18 genotyping tests for use in women with ASC-US cytology. *Obstet Gynecol Surv.* 2011;66:211-212.
171. Jovanovic AM, Dikic SD, Jovanovic V et al. Correlation of human papilloma virus infection with cytology, colposcopy and histopathological examination of the bioptic tissue in low- and high-grade intraepithelial lesions. *European Journal of Gynaecological Oncology.* 2012;33:512-516.
172. Kares S, Kujala P, Vuento R et al. HPV testing as the primary screening method in an organized regional screening program for cervical cancer. Preliminary results. *Cytopathology.* 2014;25:19.
173. Katki HA, Kinney WK, Fetterman B et al. Cervical cancer risk for women undergoing concurrent testing for human papillomavirus and cervical cytology: a population-based study in routine clinical practice.[Erratum appears in *Lancet Oncol.* 2011 Aug;12(8):722]. *Lancet Oncology.* 2011;12:663-672.
174. Katki HA, Schiffman M, Castle PE et al. Five-year risks of CIN 2+ and CIN 3+ among women with HPV-positive and HPV-negative LSIL Pap results. *J Lower Genital Tract Dis.* 2013;17:S43-S49.
175. Katki HA, Gage JC, Schiffman M et al. Follow-up testing after colposcopy: five-year risk of CIN 2+ after a colposcopic diagnosis of CIN 1 or less. *J Lower Genital Tract Dis.* 2013;17:S69-S77.
176. Katki HA, Schiffman M, Castle PE et al. Benchmarking CIN 3+ risk as the basis for incorporating HPV and Pap cotesting into cervical screening and management guidelines. *J Lower Genital Tract Dis.* 2013;17:S28-S35.
177. -----Five-year risks of CIN 3+ and cervical cancer among women with HPV testing of ASC-US Pap results. *J Low Genit Tract Dis.* 2013;17:S36-S42.
178. -----Five-year risks of CIN 3+ and cervical cancer among women who test Pap-negative but are HPV-positive. *J Low Genit Tract Dis.* 2013;17:S56-S63.
179. Kececioğlu M, Seckin B, Baser E et al. Cost and effectiveness comparison of immediate colposcopy versus human papillomavirus DNA testing in management of atypical squamous cells of undetermined significance in Turkish women. *Asian Pacific Journal of Cancer Prevention: Apjcp.* 2013;14:511-514.
180. Keegan H, Pilkington L, McInerney J et al. Human papillomavirus detection and genotyping, by HC2, full-spectrum HPV and molecular beacon real-time HPV assay in an Irish colposcopy clinic. *Journal of Virological Methods.* 2014;201:93-100.
181. Kelly RS, Patnick J, Kitchener HC et al. HPV testing as a triage for borderline or mild dyskaryosis on cervical cytology: results from the Sentinel Sites study. *British journal of cancer.* 2011;105:983-988.

182. Khan MJ, Castle PE, Lorincz AT et al. The elevated 10-year risk of cervical precancer and cancer in women with human papillomavirus (HPV) type 16 or 18 and the possible utility of type-specific HPV testing in clinical practice. *J Natl Cancer Inst.* 2005;97:1072-1079.
183. Kim JH, Kim IW, Kim YW et al. Comparison of single-, double- and triple-combined testing, including Pap test, HPV DNA test and cervicography, as screening methods for the detection of uterine cervical cancer. *Oncology Reports.* 2013;29:1645-1651.
184. Kitchener HC, Canfell K, Gilham C et al. The clinical effectiveness and cost-effectiveness of primary human papillomavirus cervical screening in England: extended follow-up of the ARTISTIC randomised trial cohort through three screening rounds. *Health Technology Assessment.* 2014;18:1-196.
185. Kitchener HC, Walker PG, Nelson L et al. HPV testing as an adjunct to cytology in the follow up of women treated for cervical intraepithelial neoplasia. *BJOG: An International Journal of Obstetrics & Gynaecology.* 2008;115:1001-1007.
186. Kitchener HC, Gilham C, Sargent A et al. A comparison of HPV DNA testing and liquid based cytology over three rounds of primary cervical screening: extended follow up in the ARTISTIC trial. *Eur J Cancer.* 2011;47:864-871.
187. Ko EM, Tambouret R, Wilbur D et al. HPV Reflex Testing in Menopausal Women. *Pathology Research International.* 2011;2011:181870.
188. Kocken M, Berkhof J, van Kemenade FJ et al. Long-term CIN3+ risk in women with abnormal cytology; role of hrHPV testing. *British journal of cancer.* 2012;106:817-825.
189. Kotaniemi-Talonen L, Nieminen P, Anttila A et al. Routine cervical screening with primary HPV testing and cytology triage protocol in a randomised setting. *British journal of cancer.* 2005;93:862-867.
190. Kotaniemi-Talonen L, Anttila A, Malila N et al. Screening with a primary human papillomavirus test does not increase detection of cervical cancer and intraepithelial neoplasia 3. *European Journal of Cancer.* 2008;44:565-571.
191. Kovacic MB, Katki HA, Kreimer AR et al. Epidemiologic analysis of histologic cervical inflammation: relationship to human papillomavirus infections. *Human Pathology.* 2008;39:1088-1095.
192. Krambeck WM, Cadide RM, Dalmarco EM et al. HPV detection and genotyping as an earlier approach in cervical cancer screening of the female genital tract. *Clinical & Experimental Obstetrics & Gynecology.* 2008;35:175-178.
193. Kroupis C, Thomopoulou G, Papatomas TG et al. Population-based study of human papillomavirus infection and cervical neoplasia in Athens, Greece. *Epidemiology & Infection.* 2007;135:943-950.
194. Kulasingam SL, Kim JJ, Lawrence WF et al. Cost-effectiveness analysis based on the atypical squamous cells of undetermined significance/low-grade squamous intraepithelial lesion Triage Study (ALTS). *Journal of the National Cancer Institute.* 2006;98:92-100.
195. Lai CH, Chao A, Chang CJ et al. Host and viral factors in relation to clearance of human papillomavirus infection: a cohort study in Taiwan. *International journal of cancer.* 2008;123:1685-1692.
196. Lazcano-Ponce E, Lorincz AT, Salmeron J et al. A pilot study of HPV DNA and cytology testing in 50,159 women in the routine Mexican Social Security Program. *Cancer Causes Control.* 2010;21:1693-1700.

197. Le DM, Giuffre G, Caruso C et al. Human papillomavirus types distribution in eastern Sicilian females with cervical lesions. A correlation with colposcopic and histological findings. *Pathology Oncology Research*. 2013;19:481-487.
198. Lee B, Suh DH, Kim K et al. Utility of Human Papillomavirus Genotyping for Triage of Patients with Atypical Squamous Cells of Undetermined Significance by Cervical Cytology. *Anticancer Research*. 2015;35:4197-4202.
199. Lee JK, Kim MK, Song SH et al. Comparison of human papillomavirus detection and typing by hybrid capture 2, linear array, DNA chip, and cycle sequencing in cervical swab samples. *Int J Gynecol Cancer*. 2009;19:266-272.
200. Lee KJ, Lee JK, Saw HS. Can human papillomavirus DNA testing substitute for cytology in the detection of high-grade cervical lesions? *Archives of Pathology & Laboratory Medicine*. 2004;128:298-302.
201. Lee SJ, Jung KL, Lee JW et al. Analyses of atypical squamous cells refined by the 2001 Bethesda System: the distribution and clinical significance of follow-up management. *Int J Gynecol Cancer*. 2006;16:664-669.
202. Leeson SC, Alibegashvili T, Arbyn M et al. HPV testing and vaccination in Europe. [Review]. *J Lower Genital Tract Dis*. 2014;18:61-69.
203. Leinonen MK, Anttila A, Malila N et al. Type- and age-specific distribution of human papillomavirus in women attending cervical cancer screening in Finland. *British journal of cancer*. 2013;109:2941-2950.
204. Li K, Yin R. Repeat cytology and human papillomavirus screening strategies in detecting preinvasive cervical lesions. *Medicine*. 2015;94:e435.
205. Liao GD, Sellors JW, Sun HK et al. p16INK4A immunohistochemical staining and predictive value for progression of cervical intraepithelial neoplasia grade 1: a prospective study in China. *International journal of cancer*. 2014;134:1715-1724.
206. Lin CJ, Lai HC, Wang KH et al. Testing for methylated PCDH10 or WT1 is superior to the HPV test in detecting severe neoplasms (CIN3 or greater) in the triage of ASC-US smear results. *American Journal of Obstetrics & Gynecology*. 2011;204:21-27.
207. Liou YL, Zhang Y, Liu Y et al. Comparison of HPV genotyping and methylated ZNF582 as triage for women with equivocal liquid-based cytology results. *Clinical Epigenetics*. 2015;7:50.
208. Litjens RJ, Theelen W, van de Pas Y et al. Use of the HPV MLPA assay in cervical cytology for the prediction of high grade lesions. *Journal of Medical Virology*. 2013;85:1386-1393.
209. Litjens RJ, Van de Vijver KK, Hopman AH et al. The majority of metachronous CIN1 and CIN3 lesions are caused by different human papillomavirus genotypes, indicating that the presence of CIN1 seems not to determine the risk for subsequent detection of CIN3. *Human Pathology*. 2014;45:221-226.
210. Littell RD, Kinney W, Fetterman B et al. Risk of cervical precancer and cancer in women aged 30 years and older with an HPV-negative low-grade squamous intraepithelial lesion screening result. *J Lower Genital Tract Dis*. 2011;15:54-59.
211. Liu TY, Xie R, Luo L et al. Diagnostic validity of human papillomavirus E6/E7 mRNA test in cervical cytological samples. *Journal of Virological Methods*. 2014;196:120-125.
212. Liverani CA, Ciavattini A, Monti E et al. High risk HPV DNA subtypes and E6/E7 mRNA expression in a cohort of colposcopy patients from Northern Italy with high-grade histologically verified cervical lesions. *Amer J Transl Res*. 2012;4:452-457.

213. Loiacono R, Falco G, Kardhashi A et al. Retrospective study of 769 patients followed at the National Cancer Institute IRCCS IΓÇ£Giovanni Paolo IIIΓÇØ, Bari (Italy). *Int J Gynecol Obstet.* 2012;119:S647.
214. Louvanto K, Syrjanen KJ, Rintala MA et al. Human papillomavirus and predictors of cervical intraepithelial neoplasia among young mothers in a prospective follow-up study. *Acta Obstetricia et Gynecologica Scandinavica.* 2011;90:167-173.
215. Louwers JA, Kocken M, van der Bijl JC et al. Colposcopic characteristics of high-risk human papillomavirus-related cervical lesions. *J Lower Genital Tract Dis.* 2010;14:49-55.
216. Luttmmer R, Berkhof J, Dijkstra MG et al. Comparing triage algorithms using HPV DNA genotyping, HPV E7 mRNA detection and cytology in high-risk HPV DNA-positive women. *J Clin Virol.* 2015;67:59-66.
217. Maenpaa J, Kares S, Kujala P et al. Hpv testing as the primary screening method in an organized regional screening program for cervical cancer. Preliminary results. *Int J Gynecol Cancer.* 2014;24:677.
218. Massad LS, Einstein MH, Huh WK et al. 2012 updated consensus guidelines for the management of abnormal cervical cancer screening tests and cancer precursors.[Erratum appears in J Low Genit Tract Dis. 2013 Jul;17(3):367]. *J Lower Genital Tract Dis.* 2013;17:Suppl-S27.
219. Massad LS. Developing guidelines for cervical cancer prevention in the face of diagnostic complexity. [Review]. *Journal of the National Comprehensive Cancer Network.* 2014;12:349-353.
220. Maucort-Boulch D, Plummer M, Castle PE et al. Predictors of human papillomavirus persistence among women with equivocal or mildly abnormal cytology. *International journal of cancer.* 2010;126:684-691.
221. Mayrand MH, Duarte-Franco E, Rodrigues I et al. Human papillomavirus DNA versus Papanicolaou screening tests for cervical cancer. *New England Journal of Medicine.* 2007;357:1579-1588.
222. McDonald AC, Tergas AI, Kuhn L et al. Distribution of Human Papillomavirus Genotypes among HIV-Positive and HIV-Negative Women in Cape Town, South Africa. *Frontiers in Oncology.* 2014;4:48.
223. McIntosh H. High-risk human papillomavirus (HPV) testing as triage for women with mild or borderline cytology abnormalities. 2012. Glasgow, NHS Quality Improvement Scotland (NHS QIS).  
Ref Type: Report
224. McKenna M, McMenamin MM. Human papillomavirus testing in young women: clinical outcomes of human papillomavirus triage in a UK cervical screening program. *Cancer Cytopathol.* 2014;122:702-710.
225. Medeiros F, Yuan L, Breslin G et al. Type-specific HPV testing as a predictor of high-grade squamous intraepithelial lesion outcome after cytologic abnormalities.[Erratum appears in J Low Genit Tract Dis. 2006 Jan;10(1):63]. *J Lower Genital Tract Dis.* 2005;9:154-159.
226. Mendoza LP, Arbiza J, Paez M et al. Distribution of human papillomavirus genotypes in Paraguayan women according to the severity of the cervical lesion. *Journal of Medical Virology.* 2011;83:1351-1357.
227. Mentzelopoulou P, Papaefthimiou M, Pouliakis A et al. The role of cytology in the management of ascus diagnosis. *Cytopathology.* 2014;25:63.



228. Menzo S, Ciavattini A, Bagnarelli P et al. Molecular epidemiology and pathogenic potential of underdiagnosed human papillomavirus types. *BMC Microbiology*. 2008;8:112.
229. Mesher D, Szarewski A, Cadman L et al. Comparison of human papillomavirus testing strategies for triage of women referred with low-grade cytological abnormalities. *European Journal of Cancer*. 2013;49:2179-2186.
230. Michelli E, Tellez L, Mendoza JA et al. Comparative analysis of three methods for HPV DNA detection in cervical samples. *Investigacion Clinica*. 2011;52:344-357.
231. Michelli E, Tellez L, Mendoza JA et al. Amplification of human papillomavirus early genes for detection of nine genotypes in Venezuelan women. *Investigacion Clinica*. 2013;54:392-405.
232. Mijit F, Ablimit T, Abduxkur G et al. Distribution of human papillomavirus (HPV) genotypes detected by routine pap smear in uyghur-muslim women from Karasay Township Hotan (Xinjiang, China). *Journal of Medical Virology*. 2015;87:1960-1965.
233. Min KJ, Yang SY, Koo YJ et al. ASC-US and LSIL triage in Korean women: Revisiting the 2012 ASCCP screening guidelines. *Gynecol Oncol*. 2015;137:72.
234. Moarcas M, Georgescu IC, Moarcas R et al. HPV-DNA testing for patients with ASC-US helps identify the women who have a high risk for precancerous cervical lesions. *Journal of Medicine & Life*. 2014;7:Spec-80.
235. Monsonego J, Pollini G, Evrard MJ et al. Linear array genotyping and hybrid capture II assay in detecting human papillomavirus genotypes in women referred for colposcopy due to abnormal Papanicolaou smear. *International Journal of STD & AIDS*. 2008;19:385-392.
236. Monsonego J, Pollini G, Evrard MJ et al. Detection of human papillomavirus genotypes among high-risk women: a comparison of hybrid capture and linear array tests. *Sex Transm Dis*. 2008;35:521-527.
237. Monsonego J, Hudgens MG, Zerat L et al. Evaluation of oncogenic human papillomavirus RNA and DNA tests with liquid-based cytology in primary cervical cancer screening: the FASE study. *International journal of cancer*. 2011;129:691-701.
238. Monsonego J, Zerat L, Syrjanen K et al. Prevalence of type-specific human papillomavirus infection among women in France: Implications for screening, vaccination, and a future generation of multivalent HPV vaccines. *Vaccine*. 2012;30:5215-5221.
239. Moore G, Fetterman B, Cox JT et al. Lessons from practice: risk of CIN 3 or cancer associated with an LSIL or HPV-positive ASC-US screening result in women aged 21 to 24. *J Lower Genital Tract Dis*. 2010;14:97-102.
240. Moss S, Gray A, Legood R et al. Effect of testing for human papillomavirus as a triage during screening for cervical cancer: observational before and after study. *BMJ*. 2006;332:83-85.
241. Moy LM, Zhao FH, Li LY et al. Human papillomavirus testing and cervical cytology in primary screening for cervical cancer among women in rural China: comparison of sensitivity, specificity, and frequency of referral. *International journal of cancer*. 2010;127:646-656.
242. Musa J, Taiwo B, Goldsmith S et al. Predictors of atypical squamous cell of undetermined significance cervical cytology with high-risk human papilloma virus genotypes. *Archives of Gynecology & Obstetrics*. 2011;283:343-348.
243. Muwonge R, Wesley RS, Nene BM et al. Evaluation of cytology and visual triage of human papillomavirus-positive women in cervical cancer prevention in India. *International journal of cancer*. 2014;134:2902-2909.

244. Nakamura Y, Matsumoto K, Satoh T et al. HPV genotyping for triage of women with abnormal cervical cancer screening results: a multicenter prospective study. *Int J Clin Oncol*. 2015.
245. Nam EJ, Kim JW, Hong JW et al. Expression of the p16 and Ki-67 in relation to the grade of cervical intraepithelial neoplasia and high-risk human papillomavirus infection. *Journal of Gynecologic Oncology*. 2008;19:162-168.
246. Namugenyi SB, Balsan MJ, Glick SN et al. Prevalence and genotype distribution of human papillomavirus in cytology specimens containing atypical glandular cells: a case-control study. *J Clin Virol*. 2013;58:432-436.
247. Nasioutziki M, Daniilidis A, Dinas K et al. The evaluation of p16INK4a immunoexpression/immunostaining and human papillomavirus DNA test in cervical liquid-based cytological samples. *Int J Gynecol Cancer*. 2011;21:79-85.
248. Naucler P, Ryd W, Tornberg S et al. HPV type-specific risks of high-grade CIN during 4 years of follow-up: a population-based prospective study. *British journal of cancer*. 2007;97:129-132.
249. Naucler P, Ryd W, Tornberg S et al. Human papillomavirus and Papanicolaou tests to screen for cervical cancer.[Erratum appears in N Engl J Med. 2008 Oct 9;359(15):1637 Note: Johansson, Bo [added]]. *New England Journal of Medicine*. 2007;357:1589-1597.
250. Naucler P, Ryd W, Tornberg S et al. Efficacy of HPV DNA testing with cytology triage and/or repeat HPV DNA testing in primary cervical cancer screening. *Journal of the National Cancer Institute*. 2009;101:88-99.
251. Ngugi CW, Schmidt D, Wanyoro K et al. p16(INK4a)/Ki-67 dual stain cytology for cervical cancer screening in Thika district, Kenya. *Infectious Agents & Cancer [Electronic Resource]*. 2015;10:25.
252. Niccolai LM, Russ C, Julian PJ et al. Individual and geographic disparities in human papillomavirus types 16/18 in high-grade cervical lesions: Associations with race, ethnicity, and poverty. *Cancer*. 2013;119:3052-3058.
253. Nomelini RS, Barcelos AC, Michelin MA et al. Utilization of human papillomavirus testing for cervical cancer prevention in a university hospital. *Cadernos de Saude Publica*. 2007;23:1309-1318.
254. Nomelini RS, Guimaraes PD, Candido PA et al. Prevention of cervical cancer in women with ASCUS in the Brazilian Unified National Health System: cost-effectiveness of the molecular biology method for HPV detection. *Cadernos de Saude Publica*. 2012;28:2043-2052.
255. Nygard M, Hansen BT, Dillner J et al. Targeting human papillomavirus to reduce the burden of cervical, vulvar and vaginal cancer and pre-invasive neoplasia: establishing the baseline for surveillance. *PLoS ONE [Electronic Resource]*. 2014;9:e88323.
256. O'Connor C, Myles H, O'Connor MB et al. An audit of Colposcopy referrals from a GU/STD clinic. *BMC Research Notes*. 2008;1:24.
257. Ogilvie GS, van Niekerk DJ, Kraiden M et al. A randomized controlled trial of Human Papillomavirus (HPV) testing for cervical cancer screening: trial design and preliminary results (HPV FOCAL Trial). *BMC cancer*. 2010;Vol.10, pp.111, 2010..
258. Ogilvie GS, Kraiden M, van Niekerk DJ et al. Primary cervical cancer screening with HPV testing compared with liquid-based cytology: results of round 1 of a randomised controlled trial -- the HPV FOCAL Study. *British journal of cancer*. 2012;107:1917-1924.

259. Okadome M, Saito T, Tanaka H et al. Potential impact of combined high- and low-risk human papillomavirus infection on the progression of cervical intraepithelial neoplasia 2. *Journal of Obstetrics & Gynaecology Research*. 2014;40:561-569.
260. Ovestad IT, Vennestrom U, Andersen L et al. Comparison of different commercial methods for HPV detection in follow-up cytology after ASCUS/LSIL, prediction of CIN2-3 in follow up biopsies and spontaneous regression of CIN2-3. *Gynecol Oncol*. 2011;123:278-283.
261. Paba P, Ascone C, Criscuolo AA et al. Human papillomavirus molecular testing in women with low grade cervical lesions: experience from an Italian hospital. *Anticancer Research*. 2012;32:1253-1257.
262. Pacchiarotti A, Ferrari F, Bellardini P et al. Prognostic value of p16-INK4A protein in women with negative or CIN1 histology result: a follow-up study. *International journal of cancer*. 2014;134:897-904.
263. Padalko E, Van RL, Bamelis M et al. Prospective evaluation of E6/E7 mRNA detection by the NucliSENS Easy Q HPV assay in a stepwise protocol. *Journal of Medical Virology*. 2013;85:1242-1249.
264. Paengchit K, Kietpeerakool C, Wangchai W et al. Cervical pathology in cytology-negative/HPV-positive women: results from Lampang Cancer Hospital, Thailand. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2014;15:7951-7954.
265. Pajtler M, Milicic-Juhas V, Milojkovic M et al. Assessment of HPV DNA test value in management women with cytological findings of ASC-US, CIN1 and CIN2. *Collegium Antropologicum*. 2010;34:81-86.
266. Pan QJ, Hu SY, Zhang X et al. Pooled analysis of the performance of liquid-based cytology in population-based cervical cancer screening studies in China. *Cancer Cytopathol*. 2013;121:473-482.
267. Pan QJ, Hu SY, Guo HQ et al. Liquid-based cytology and human papillomavirus testing: a pooled analysis using the data from 13 population-based cervical cancer screening studies from China. *Gynecol Oncol*. 2014;133:172-179.
268. Pasquale L, Giorgi RP, Carozzi F et al. Cervical cancer screening with HPV testing in the Valcamonica (Italy) screening programme. *Journal of Medical Screening*. 2015;22:38-48.
269. Perksanusak T, Sananpanichkul P, Chirdchim W et al. Colposcopy Requirement of Papanicolaou Smear after Atypical Squamous Cells of Undetermined Significance (ASC-US) by Follow-up Protocol in an Urban Gynaecology Clinic, a Retrospective Study in Thailand. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2015;16:4977-4980.
270. Perrons C, Jelley R, Kleter B et al. Detection of persistent high risk human papillomavirus infections with hybrid capture II and SPF10/LiPA. *J Clin Virol*. 2005;32:278-285.
271. Petry KU, Schmidt D, Scherbring S et al. Triaging Pap cytology negative, HPV positive cervical cancer screening results with p16/Ki-67 Dual-stained cytology. *Gynecol Oncol*. 2011;121:505-509.
272. Petry KU, Luyten A, Justus A et al. Prevalence of low-risk HPV types and genital warts in women born 1988/89 or 1983/84 -results of WOLVES, a population-based epidemiological study in Wolfsburg, Germany. *BMC Infectious Diseases*. 2012;12:367.
273. Petry KU, Luyten A, Scherbring S. Accuracy of colposcopy management to detect CIN3 and invasive cancer in women with abnormal screening tests: results from a primary HPV screening project from 2006 to 2011 in Wolfsburg, Germany. *Gynecol Oncol*. 2013;128:282-287.

274. Piccoli R, Mandato VD, Lavitola G et al. Atypical squamous cells and low squamous intraepithelial lesions in postmenopausal women: implications for management. *European Journal of Obstetrics, Gynecology, & Reproductive Biology*. 2008;140:269-274.
275. Pity IS, Shamdeen MY, Wais SA. Follow up of atypical squamous cell Pap smears in Iraqi women. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2012;13:3455-3460.
276. Possati-Resende JC, Fregnani JH, Kerr LM et al. The Accuracy of p16/Ki-67 and HPV Test in the Detection of CIN2/3 in Women Diagnosed with ASC-US or LSIL. *PLoS ONE [Electronic Resource]*. 2015;10:e0134445.
277. Pouliakis A, Karakitsou E, Chrelias C et al. The Application of Classification and Regression Trees for the Triage of Women for Referral to Colposcopy and the Estimation of Risk for Cervical Intraepithelial Neoplasia: A Study Based on 1625 Cases with Incomplete Data from Molecular Tests. *BioMed Research International*. 2015;2015:914740.
278. Pretorius RG, Belinson JL, Peterson P et al. Yield and mode of diagnosis of cervical intraepithelial neoplasia 3 or cancer among women with negative cervical cytology and positive high-risk human papillomavirus test results. *J Lower Genital Tract Dis*. 2013;17:430-439.
279. Pruski D, Fraszczak J, Iwaniec K et al. Assessment of frequency of regression and progression of mild cervical neoplasia--LGSIL in women with positive high-risk HPV DNA test result. *Ginekologia Polska*. 2012;83:572-575.
280. Qiao YL, Jeronimo J, Zhao FH et al. Lower cost strategies for triage of human papillomavirus DNA-positive women. *International journal of cancer*. 2014;134:2891-2901.
281. Rabelo-Santos SH, Derchain SF, Villa LL et al. Human papillomavirus-specific genotypes in cervical lesions of women referred for smears with atypical glandular cells or adenocarcinoma in situ. *International Journal of Gynecological Pathology*. 2009;28:272-278.
282. Ramzan M, Ain NU, Ilyas S et al. A cornucopia of screening and diagnostic techniques for human papillomavirus associated cervical carcinomas. [Review]. *Journal of Virological Methods*. 2015;222:192-201.
283. Rao A, Sandri MT, Sideri M et al. Comparison of hybrid capture 2 High-Risk HPV results in the low positive range with cobas HPV Test results from the ATHENA study. *J Clin Virol*. 2013;58:161-167.
284. Ratnam S, Coutlee F, Fontaine D et al. Clinical performance of the PreTect HPV-Proofer E6/E7 mRNA assay in comparison with that of the Hybrid Capture 2 test for identification of women at risk of cervical cancer. *J Clin Microbiol*. 2010;48:2779-2785.
285. ----Aptima HPV E6/E7 mRNA test is as sensitive as Hybrid Capture 2 Assay but more specific at detecting cervical precancer and cancer. *J Clin Microbiol*. 2011;49:557-564.
286. Rebolj M, Lynge E, Bonde J. Human papillomavirus testing and genotyping in cervical screening. [Review]. *Expert Review of Anticancer Therapy*. 2011;11:1023-1031.
287. Rebolj M, Lynge E, Ejegod D et al. Comparison of three human papillomavirus DNA assays and one mRNA assay in women with abnormal cytology. *Gynecol Oncol*. 2014;135:474-480.
288. Rebolj M, Bonde J, Ejegod D et al. A daunting challenge: Human Papillomavirus assays and cytology in primary cervical screening of women below age 30years. *European Journal of Cancer*. 2015;51:1456-1466.
289. Reid JL, Wright TC, Jr., Stoler MH et al. Human Papillomavirus Oncogenic mRNA Testing for Cervical Cancer Screening: Baseline and Longitudinal Results From the CLEAR Study. *American Journal of Clinical Pathology*. 2015;144:473-483.

290. Riera BM, Ruperez PB, L+ízaró V, I et al. High risk HPV, reason of screening and progression. *Int J Gynecol Cancer*. 2013;23:781.
291. Rijkaart DC, Berkhof J, van Kemenade FJ et al. Evaluation of 14 triage strategies for HPV DNA-positive women in population-based cervical screening. *International journal of cancer*. 2012;130:602-610.
292. Rijkaart DC, Heideman DA, Coupe VM et al. High-risk human papillomavirus (hrHPV) E6/E7 mRNA testing by PreTect HPV-Proofer for detection of cervical high-grade intraepithelial neoplasia and cancer among hrHPV DNA-positive women with normal cytology. *J Clin Microbiol*. 2012;50:2390-2396.
293. Rijkaart DC, Berkhof J, van Kemenade FJ et al. HPV DNA testing in population-based cervical screening (VUSA-Screen study): results and implications. *British journal of cancer*. 2012;106:975-981.
294. Rodriguez AC, Schiffman M, Herrero R et al. Rapid clearance of human papillomavirus and implications for clinical focus on persistent infections. *Journal of the National Cancer Institute*. 2008;100:513-517.
295. Rodriguez AC, Schiffman M, Herrero R et al. Longitudinal study of human papillomavirus persistence and cervical intraepithelial neoplasia grade 2/3: critical role of duration of infection. *Journal of the National Cancer Institute*. 2010;102:315-324.
296. Rokita W, Kedzia W, Pruski D et al. Comparison of the effectiveness of cytodiagnosics, molecular identification of HPV HR and CINtecPLUS test to identify LG SIL and HG SIL. *Ginekologia Polska*. 2012;83:894-898.
297. Ronco G, Giorgi-Rossi P, Carozzi F et al. Efficacy of human papillomavirus testing for the detection of invasive cervical cancers and cervical intraepithelial neoplasia: a randomised controlled trial. *Lancet Oncology*. 2010;11:249-257.
298. Ronco G, Dillner J, Elfstrom KM et al. Efficacy of HPV-based screening for prevention of invasive cervical cancer: follow-up of four European randomised controlled trials. *Lancet*. 2014;383:524-532.
299. Saccardi C, Gizzo S, Noventa M et al. High-risk human papillomavirus DNA test: could it be useful in low-grade cervical lesion triage? Five-year follow-up. *Reprod Sci*. 2014;21:198-203.
300. Safaeian M, Schiffman M, Gage J et al. Detection of precancerous cervical lesions is differential by human papillomavirus type. *Cancer Research*. 2009;69:3262-3266.
301. Sahasrabudde VV, Gravitt PE, Dunn ST et al. Comparison of human papillomavirus detections in urine, vulvar, and cervical samples from women attending a colposcopy clinic. *J Clin Microbiol*. 2014;52:187-192.
302. Sahasrabudde VV, Gravitt PE, Dunn ST et al. Evaluation of clinical performance of a novel urine-based HPV detection assay among women attending a colposcopy clinic. *J Clin Virol*. 2014;60:414-417.
303. Samarawardana P, Dehn DL, Singh M et al. p16(INK4a) is superior to high-risk human papillomavirus testing in cervical cytology for the prediction of underlying high-grade dysplasia.[Erratum appears in *Cancer Cytopathol*. 2010 Oct 25;118(5):303]. *Cancer Cytopathol*. 2010;118:146-156.
304. Sankaranarayanan R, Nene BM, Shastri SS et al. HPV screening for cervical cancer in rural India. *New England Journal of Medicine*. 2009;360:1385-1394.
305. Saslow D, Solomon D, Lawson HW et al. American Cancer Society, American Society for Colposcopy and Cervical Pathology, and American Society for Clinical Pathology screening

- guidelines for the prevention and early detection of cervical cancer. *CA: a Cancer Journal for Clinicians*. 2012;62:147-172.
306. Sauer T, Morland E, Lie KA et al. Results from the first 5 years of secondary HPV testing in the norwegian cervical cancer screening program. *Cytopathology*. 2011;22:47-48.
  307. Sauter JL, Mount SL, St John TL et al. Testing of integrated human papillomavirus mRNA decreases colposcopy referrals: could a change in human papillomavirus detection methodology lead to more cost-effective patient care? *Acta Cytologica*. 2014;58:162-166.
  308. Schettino MT, Ammaturo FP, Grimaldi E et al. Persistent papillomavirus type-31 and type-45 infections predict the progression to squamous intraepithelial lesion. *Taiwanese Journal of Obstetrics & Gynecology*. 2014;53:494-497.
  309. Schiffman M, Wentzensen N, Wacholder S et al. Human papillomavirus testing in the prevention of cervical cancer. [Review]. *Journal of the National Cancer Institute*. 2011;103:368-383.
  310. Schiffman M, Glass AG, Wentzensen N et al. A long-term prospective study of type-specific human papillomavirus infection and risk of cervical neoplasia among 20,000 women in the Portland Kaiser Cohort Study. *Cancer Epidemiol Biomarkers Prev*. 2011;20:1398-1409.
  311. Schiffman M, Vaughan LM, Raine-Bennett TR et al. A study of HPV typing for the management of HPV-positive ASC-US cervical cytologic results. *Gynecol Oncol*. 2015;138:573-578.
  312. Schiffman M, Boyle S, Raine-Bennett T et al. The Role of Human Papillomavirus Genotyping in Cervical Cancer Screening: A Large-Scale Evaluation of the cobas HPV Test. *Cancer Epidemiology, Biomarkers & Prevention*. 2015;24:1304-1310.
  313. Schiffman M, Burk RD, Boyle S et al. A study of genotyping for management of human papillomavirus-positive, cytology-negative cervical screening results. *J Clin Microbiol*. 2015;53:52-59.
  314. Schlichte MJ, Guidry J. Current Cervical Carcinoma Screening Guidelines. [Review]. *Journal of Clinical Medicine*. 2015;4:918-932.
  315. Schnatz PF, Sharpless KE, O'Sullivan DM. Use of human papillomavirus testing in the management of atypical glandular cells. *J Lower Genital Tract Dis*. 2009;13:94-101.
  316. Sellors JW, Schweizer JG, Lu PS et al. Association of elevated E6 oncoprotein with grade of cervical neoplasia using PDZ interaction-mediated precipitation of E6. *J Lower Genital Tract Dis*. 2011;15:169-176.
  317. Selvaggi SM. The utility of high-risk HPV testing in the management of women 30 to 64 years of age with low-grade squamous intraepithelial lesions on cervical cytology. *J Am Soc Cytopathology*. 2015;4:290-293.
  318. Seo SS, Song YS, Kim JW et al. Good correlation of HPV DNA test between self-collected vaginal and clinician-collected cervical samples by the oligonucleotide microarray. *Gynecol Oncol*. 2006;102:67-73.
  319. Shahid M, Kazmi SU, Rehman A et al. Cervical cancer screening and HPV genotype distribution among asymptomatic patients of Karachi Pakistan. *Pakistan Journal of Medical Sciences*. 2015;31:493-498.
  320. Shastri SS, Dinshaw K, Amin G et al. Concurrent evaluation of visual, cytological and HPV testing as screening methods for the early detection of cervical neoplasia in Mumbai, India. *Bulletin of the World Health Organization*. 2005;83:186-194.

321. Shen-Gunther J, Yu X. HPV molecular assays: defining analytical and clinical performance characteristics for cervical cytology specimens. *Gynecol Oncol*. 2011;123:263-271.
322. Shen Y, Gong J, He Y et al. Quantivirus HPV E6/E7 RNA 3.0 assay (bDNA) is as sensitive, but less specific than Hybrid Capture 2 test.[Erratum appears in J Virol Methods. 2013 Nov;193(2):729]. *Journal of Virological Methods*. 2013;187:288-293.
323. Shi JF, Belinson JL, Zhao FH et al. Human papillomavirus testing for cervical cancer screening: results from a 6-year prospective study in rural China. *American Journal of Epidemiology*. 2009;170:708-716.
324. Shipitsyna E, Zolotoverkhaya E, Kuevda D et al. Prevalence of high-risk human papillomavirus types and cervical squamous intraepithelial lesions in women over 30 years of age in St. Petersburg, Russia. *Cancer Epidemiology*. 2011;35:160-164.
325. Shwe MM, Kyi KN, Okada S et al. Trend of human papillomavirus genotypes in cervical neoplasia observed in a newly developing township in Yangon, Myanmar. *Acta Medica Okayama*. 2015;69:51-58.
326. Siebers AG, Arbyn M, Melchers WJG et al. Effectiveness of two strategies to follow-up ASC-US and LSIL screening results in The Netherlands using repeat cytology with or without additional hrHPV testing: a retrospective cohort study. *Cancer Causes Control*. 2014.
327. Sigurdsson K, Sigvaldason H, Gudmundsdottir T et al. The efficacy of HPV 16/18 vaccines on sexually active 18-23 year old women and the impact of HPV vaccination on organized cervical cancer screening. *Acta Obstetrica et Gynecologica Scandinavica*. 2009;88:27-35.
328. Sikon AL, Jin XW, Lipold LD et al. Pap vs. HPV vs co-test for CIN2+ detection: A retrospective study of performance and cost. *J Gen Intern Med*. 2014;29:S163-S164.
329. Silva C, Almeida EC, Cobo EC et al. A retrospective study on cervical intraepithelial lesions of low-grade and undetermined significance: evolution, associated factors and cytohistological correlation. *Sao Paulo Medical Journal = Revista Paulista de Medicina*. 2014;132:92-96.
330. Sodhani P, Gupta S, Sharma JK et al. Test characteristics of various screening modalities for cervical cancer: a feasibility study to develop an alternative strategy for resource-limited settings. *Cytopathology*. 2006;17:348-352.
331. Sole-Sedeno JM, Mancebo G, Miralpeix E et al. Utility of HPV genotyping in the management of low-grade squamous intraepithelial cervical lesions (LSIL). *Gynecol Oncol*. 2015;137:31.
332. Solomon J, Teschendorf C, Hasteh F. Management of cytology negative high-risk human papillomavirus positive cervical smears. *Lab Invest*. 2015;95:308A-309A.
333. Song SH, Lee JK, Oh MJ et al. Risk factors for the progression or persistence of untreated mild dysplasia of the uterine cervix. *Int J Gynecol Cancer*. 2006;16:1608-1613.
334. Sorbye SW, Arbyn M, Fismen S et al. HPV E6/E7 mRNA testing is more specific than cytology in post-colposcopy follow-up of women with negative cervical biopsy. *PLoS ONE [Electronic Resource]*. 2011;6:e26022.
335. Sorbye SW, Fismen S, Gutteberg TJ et al. HPV mRNA testing in triage of women with ASC-US cytology may reduce the time for CIN2+diagnosis compared with repeat cytology. *Current Pharmaceutical Design*. 2013;19:1401-1405.
336. -----HPV mRNA is more specific than HPV DNA in triage of women with minor cervical lesions. *PLoS ONE [Electronic Resource]*. 2014;9:e112934.

337. Soto Y, Torres G, Kouri V et al. Molecular epidemiology of human papillomavirus infections in cervical samples from Cuban women older than 30 years. *J Lower Genital Tract Dis.* 2014;18:210-217.
338. Spathis A, Kottaridi C, Chranioti A et al. mRNA and DNA detection of human papillomaviruses in women of all ages attending two colposcopy clinics. *PLoS ONE [Electronic Resource].* 2012;7:e49205.
339. Spinillo A, Dal BB, Gardella B et al. Multiple human papillomavirus infection and high grade cervical intraepithelial neoplasia among women with cytological diagnosis of atypical squamous cells of undetermined significance or low grade squamous intraepithelial lesions. *Gynecol Oncol.* 2009;113:115-119.
340. Spinillo A, Gardella B, Chiesa A et al. Diagnostic accuracy of colposcopy in relation to human papillomavirus genotypes and multiple infection. *Gynecol Oncol.* 2014;134:527-533.
341. Spinillo A, Gardella B, Roccio M et al. Multiple human papillomavirus infection with or without type 16 and risk of cervical intraepithelial neoplasia among women with cervical cytological abnormalities. *Cancer Causes Control.* 2014;25:1669-1676.
342. Stanculescu R, Bratila E, Bausic V et al. The triage of low-grade cytological abnormalities by the immunocytochemical expression of cyclin-dependent kinase inhibitor p16INK4a versus Human Papillomavirus test: a real possibility to predict cervical intraepithelial neoplasia CIN2 or CIN2+. *Romanian Journal of Morphology & Embryology.* 2013;54:1061-1065.
343. Stanczuk GA, Currie H, Baxter G et al. Cobas 4800 HPV detection in the cervical, vaginal and urine samples of women with high-grade CIN before and after treatment. *Journal of Clinical Pathology.* 2015;68:567-570.
344. Stoler MH, Wright TC, Jr., Sharma A et al. High-risk human papillomavirus testing in women with ASC-US cytology: results from the ATHENA HPV study. *American Journal of Clinical Pathology.* 2011;135:468-475.
345. Stoler MH, Wright TC, Jr., Cuzick J et al. APTIMA HPV assay performance in women with atypical squamous cells of undetermined significance cytology results. *American Journal of Obstetrics & Gynecology.* 2013;208:144-148.
346. Stoler MH, Austin RM, Zhao C. Point-Counterpoint: Cervical Cancer Screening Should Be Done by Primary Human Papillomavirus Testing with Genotyping and Reflex Cytology for Women over the Age of 25 Years. *J Clin Microbiol.* 2015;53:2798-2804.
347. Sultana F, Gertig DM, Wrede CD et al. A pilot study to compare dry cervical sample collection with standard practice of wet cervical samples for human papillomavirus testing. *J Clin Virol.* 2015;69:210-213.
348. Supho B, Supoken A, Kleebkaew P et al. Cervical pathology in high-risk human papillomavirus- positive, cytologically normal women. *Asian Pacific Journal of Cancer Prevention: Apjcp.* 2014;15:7977-7980.
349. Sycuro LK, Xi LF, Hughes JP et al. Persistence of genital human papillomavirus infection in a long-term follow-up study of female university students. *Journal of infectious diseases.* 2008;198:971-978.
350. Szarewski A, Cadman L, Mallett S et al. Human papillomavirus testing by self-sampling: assessment of accuracy in an unsupervised clinical setting. *Journal of Medical Screening.* 2007;14:34-42.
351. Szarewski A, Ambroisine L, Cadman L et al. Comparison of predictors for high-grade cervical intraepithelial neoplasia in women with abnormal smears. *Cancer Epidemiology, Biomarkers & Prevention.* 2008;17:3033-3042.



352. Szarewski A. Ushering in a new era: human papillomavirus (HPV) testing comes to the NHS Cervical Screening Programme. *Journal of Family Planning & Reproductive Health Care*. 2011;37:64-67.
353. Szarewski A, Mesher D, Cadman L et al. Comparison of seven tests for high-grade cervical intraepithelial neoplasia in women with abnormal smears: the Predictors 2 study. *J Clin Microbiol*. 2012;50:1867-1873.
354. Tempfer C, Grimm C, Harwanegg C et al. Frequency of 23 human papillomavirus types using DNA microarray in women with and without cytological anomalies. *Anticancer Research*. 2007;27:1721-1726.
355. Tinelli A, Guido M, Zizza A et al. The mRNA-HPV test utilization in the follow up of HPV related cervical lesions. *Current Pharmaceutical Design*. 2013;19:1458-1465.
356. Tokmak A, Guzel AI, Ozgu E et al. Clinical significance of atypical squamous cells of undetermined significance in detecting preinvasive cervical lesions in post- menopausal Turkish women. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2014;15:6639-6641.
357. Tornesello ML, Duraturo ML, Buonaguro L et al. Prevalence of human papillomavirus genotypes and their variants in high risk West Africa women immigrants in South Italy. *Infectious Agents & Cancer [Electronic Resource]*. 2007;2:1.
358. Tornesello ML, Cassese R, De RN et al. High prevalence of human papillomavirus infection in Eastern European and West African women immigrants in South Italy. *APMIS*. 2011;119:701-709.
359. Trimble CL, Piantadosi S, Gravitt P et al. Spontaneous regression of high-grade cervical dysplasia: effects of human papillomavirus type and HLA phenotype. *Clinical Cancer Research*. 2005;11:4717-4723.
360. Trope A, Sjoborg KD, Nygard M et al. Cytology and human papillomavirus testing 6 to 12 months after ASCUS or LSIL cytology in organized screening to predict high-grade cervical neoplasia between screening rounds. *J Clin Microbiol*. 2012;50:1927-1935.
361. Tsai HT, Tsai YM, Yang SF et al. A notable accessory screening program for detection of cervical intraepithelial neoplasia. *Pathologie Biologie*. 2009;57:477-482.
362. Tsiodras S, Georgoulakis J, Chranioti A et al. Hybrid capture vs. PCR screening of cervical human papilloma virus infections. Cytological and histological associations in 1270 women. *BMC cancer*. 2010;10:53.
363. Tsiodras S, Alepaki M, Chranioti A et al. NASBA in correlation with clinical array hpv testing and associations with cytological and histological diagnoses in 515 women. *Cytopathology*. 2011;22:59.
364. Tsoumpou I, Valasoulis G, Founta C et al. High-risk human papillomavirus DNA test and p16(INK4a) in the triage of LSIL: a prospective diagnostic study. *Gynecol Oncol*. 2011;121:49-53.
365. Twiggs LB, Hopkins M. High-risk HPV DNA testing and HPV-16/18 genotyping: what is the clinical application?. [Review]. *J Lower Genital Tract Dis*. 2011;15:224-230.
366. Twu NF, Yen MS, Lau HY et al. Type-specific human papillomavirus DNA testing with the genotyping array: a comparison of cervical and vaginal sampling. *European Journal of Obstetrics, Gynecology, & Reproductive Biology*. 2011;156:96-100.
367. Uijterwaal MH, Witte BI, van Kemenade FJ et al. Triage of borderline/mild dyskaryotic Pap cytology with p16/Ki-67 dual-stained cytology testing: cross-sectional and longitudinal outcome study. *British journal of cancer*. 2014;110:1579-1586.

368. Uijterwaal MH, Polman NJ, Witte BI et al. Triaging HPV-positive women with normal cytology by p16/Ki-67 dual-stained cytology testing: baseline and longitudinal data. *International journal of cancer*. 2015;136:2361-2368.
369. Ursu RG, Onofriescu M, Nemescu D et al. HPV prevalence and type distribution in women with or without cervical lesions in the Northeast region of Romania. *Virology Journal*. 2011;8:558.
370. van der Marel J, van BR, Quint WG et al. The impact of human papillomavirus genotype on colposcopic appearance: a cross-sectional analysis. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2014;121:1117-1126.
371. van der Marel J, Berkhof J, Ordi J et al. Attributing oncogenic human papillomavirus genotypes to high-grade cervical neoplasia: which type causes the lesion? *American Journal of Surgical Pathology*. 2015;39:496-504.
372. Van Le L. Summary of "Primary cervical cancer screening with human papillomavirus: End of study results from the ATHENA study using HPV as the first-line screening test" by Wright et al 2015. *Obstet Gynecol Surv*. 2015;70:321-322.
373. Verdoodt F, Szarewski A, Halfon P et al. Triage of women with minor abnormal cervical cytology: meta-analysis of the accuracy of an assay targeting messenger ribonucleic acid of 5 high-risk human papillomavirus types. [Review]. *Cancer Cytopathol*. 2013;121:675-687.
374. Vidal AC, Smith JS, Iversen E et al. HPV genotype distribution and cervical intraepithelial neoplasia in African American and white women living in the Southeastern United States. *Cancer Prev Res*. 2013;6.
375. Vidal AC, Smith JS, Valea F et al. HPV genotypes and cervical intraepithelial neoplasia in a multiethnic cohort in the southeastern USA. *Cancer Causes Control*. 2014;25:1055-1062.
376. Vink MA, Bogaards JA, Meijer CJ et al. Primary human papillomavirus DNA screening for cervical cancer prevention: Can the screening interval be safely extended? *International journal of cancer*. 2015;137:420-427.
377. Wahlstrom C, Iftner T, Dillner J et al. Population-based study of screening test performance indices of three human papillomavirus DNA tests. *Journal of Medical Virology*. 2007;79:1169-1175.
378. Walavalkar V, Fischer AH, Owens CL. Significance of cytopathologist's review of Pap tests screened as negative for intraepithelial lesion or malignancy that are positive for high-risk human papillomavirus. *J Am Soc Cytopathology*. 2015;4:190-194.
379. Walker JL, Wang SS, Schiffman M et al. Predicting absolute risk of CIN3 during post-colposcopic follow-up: results from the ASCUS-LSIL Triage Study (ALTS). *American Journal of Obstetrics & Gynecology*. 2006;195:341-348.
380. Walmer DK, Eder PS, Bell L et al. Human papillomavirus prevalence in a population of women living in Port-au-Prince and Leogane, Haiti. *PLoS ONE [Electronic Resource]*. 2013;8:e76110.
381. Wang JL, Yang YZ, Dong WW et al. Application of human papillomavirus in screening for cervical cancer and precancerous lesions. *Asian Pacific Journal of Cancer Prevention: Apjcp*. 2013;14:2979-2982.
382. Wang SS, Zuna RE, Wentzensen N et al. Human papillomavirus cofactors by disease progression and human papillomavirus types in the study to understand cervical cancer early endpoints and determinants. *Cancer Epidemiology, Biomarkers & Prevention*. 2009;18:113-120.

383. Watson M, Benard V, Lin L et al. Provider management of equivocal cervical cancer screening results among underserved women, 2009-2011: follow-up of atypical squamous cells of undetermined significance.[Erratum appears in *Cancer Causes Control*. 2015 May;26(5):765; PMID: 25924587]. *Cancer Causes Control*. 2015;26:759-764.
384. Wei H, Wang N, Zhang Y et al. Distribution of various types of low-risk human papillomavirus according to cervical cytology and histology in northern Chinese women. *International Journal of Gynaecology & Obstetrics*. 2014;126:28-32.
385. Wentzensen N, Schiffman M, Dunn T et al. Multiple human papillomavirus genotype infections in cervical cancer progression in the study to understand cervical cancer early endpoints and determinants. *International journal of cancer*. 2009;125:2151-2158.
386. Wentzensen N, Schiffman M, Dunn ST et al. Grading the severity of cervical neoplasia based on combined histopathology, cytopathology, and HPV genotype distribution among 1,700 women referred to colposcopy in Oklahoma. *International journal of cancer*. 2009;124:964-969.
387. Wentzensen N, Wilson LE, Wheeler CM et al. Hierarchical clustering of human papilloma virus genotype patterns in the ASCUS-LSIL triage study. *Cancer Research*. 2010;70:8578-8586.
388. Wentzensen N, Fetterman B, Castle PE et al. p16/Ki-67 Dual Stain Cytology for Detection of Cervical Precancer in HPV-Positive Women. *Journal of the National Cancer Institute*. 2015;107:2015.
389. Werner CL, Griffith WF, III, Ashfaq R et al. Comparison of human papilloma virus testing and spectroscopy combined with cervical cytology for the detection of high-grade cervical neoplasia. *J Lower Genital Tract Dis*. 2007;11:73-79.
390. Wheeler CM, Hunt WC, Schiffman M et al. Human papillomavirus genotypes and the cumulative 2-year risk of cervical precancer. *Journal of infectious diseases*. 2006;194:1291-1299.
391. Wheeler CM, Hunt WC, Joste NE et al. Human papillomavirus genotype distributions: implications for vaccination and cancer screening in the United States. *Journal of the National Cancer Institute*. 2009;101:475-487.
392. Wheeler CM, Hunt WC, Cuzick J et al. The influence of type-specific human papillomavirus infections on the detection of cervical precancer and cancer: A population-based study of opportunistic cervical screening in the United States. *International journal of cancer*. 2014;135:624-634.
393. White C, Keegan H, Pilkington L et al. Evaluation of the clinical performance of the cobas 4800 HPV test in patients referred for colposcopy. *J Clin Microbiol*. 2013;51:3415-3417.
394. Whitlock EP, Vesco KK, Eder M et al. Liquid-based cytology and human papillomavirus testing to screen for cervical cancer: a systematic review for the U.S. Preventive Services Task Force. [Review][Erratum appears in *Ann Intern Med*. 2012 Jan 3;156(1 Pt 1):71-2]. *Annals of Internal Medicine*. 2011;155:687-697.
395. Winer RL, Kiviat NB, Hughes JP et al. Development and duration of human papillomavirus lesions, after initial infection. *Journal of infectious diseases*. 2005;191:731-738.
396. Winsley E, Maharaj D, Abels P et al. Outcomes in HrHPV-positive women with low grade cervical smears and normal or low grade initial colposcopy results. *New Zealand Medical Journal*. 2014;127:37-43.

397. Wong AK, Chan RC, Nichols WS et al. Human papillomavirus (HPV) in atypical squamous cervical cytology: the Invader HPV test as a new screening assay. *J Clin Microbiol.* 2008;46:869-875.
398. Woodard A, Austin RM, Li Z et al. Prevalence of HPV 16/18 genotypes and histopathologic follow-up outcomes in women with negative cytology and positive high-risk HPV test results. *J Am Soc Cytopathology.* 2015;4:261-266.
399. Woodard AH, Li Z, Austin M et al. Histopathologic follow-up outcomes and prevalence of HPV 16/18 genotypes in women with negative cytology and positive high risk HPV test results in a large academic womens hospital laboratory. *Lab Invest.* 2014;94:127A.
400. Wright JD, Rader JS, Davila R et al. Human papillomavirus triage for young women with atypical squamous cells of undetermined significance. *Obstetrics & Gynecology.* 2006;107:822-829.
401. Wright T, Castle P, Behrens C et al. Risk of p16<sup>INK4</sup> after 3 years of follow-up of 42,209 screened women: End of study results from ATHENA. *Gynecol Oncol.* 2013;130:e166.
402. Wright TC, Stoler MH, Behrens CM et al. Comparison of the performance of human papillomavirus (HPV) primary screening strategies with cytology-based strategies: Results from the ATHENA trial 3-year follow-up phase. *Gynecol Oncol.* 2014;133:3.
403. Wright TC, Stoler MH, Behrens CM et al. Primary cervical cancer screening with human papillomavirus: end of study results from the ATHENA study using HPV as the first-line screening test. *Gynecol Oncol.* 2015;136:189-197.
404. Wright TC, Jr., Massad LS, Dunton CJ et al. 2006 consensus guidelines for the management of women with abnormal cervical cancer screening tests. [Review] [81 refs]. *American Journal of Obstetrics & Gynecology.* 2007;197:346-355.
405. ----2006 consensus guidelines for the management of women with cervical intraepithelial neoplasia or adenocarcinoma in situ. [90 refs][Erratum appears in J Low Genit Tract Dis. 2008 Jan;12(1):63]. *J Lower Genital Tract Dis.* 2007;11:223-239.
406. Wright TC, Jr., Stoler MH, Sharma A et al. Evaluation of HPV-16 and HPV-18 genotyping for the triage of women with high-risk HPV+ cytology-negative results. *American Journal of Clinical Pathology.* 2011;136:578-586.
407. Wright TC, Jr., Stoler MH, Behrens CM et al. Interlaboratory variation in the performance of liquid-based cytology: insights from the ATHENA trial. *International journal of cancer.* 2014;134:1835-1843.
408. Wu D, Cai L, Huang M et al. Prevalence of genital human papillomavirus infection and genotypes among women from Fujian province, PR China. *European Journal of Obstetrics, Gynecology, & Reproductive Biology.* 2010;151:86-90.
409. Wu R, Du H, Belinson SE et al. Secondary screening after primary self-sampling for human papillomavirus from SHENCCAST II. *J Lower Genital Tract Dis.* 2012;16:416-420.
410. Wu S, Meng L, Wang S et al. A comparison of four screening methods for cervical neoplasia. *International Journal of Gynaecology & Obstetrics.* 2005;91:189-193.
411. Yao G. Cervical cytology ASCUS patients with HPV detection and clinical value. *Int J Gynecol Cancer.* 2014;24:716.
412. Ye J, Cheng X, Chen X et al. Short-term type-specific HPV persistence and its predictors in an asymptomatic general female population in Zhejiang, China. *International Journal of Gynaecology & Obstetrics.* 2010;110:217-222.

413. Yetimalar H, Kasap B, Cukurova K et al. Cofactors in human papillomavirus infection and cervical carcinogenesis. *Archives of Gynecology & Obstetrics*. 2012;285:805-810.
414. You K, Guo YL, Geng L et al. Analysis of one year follow-up of women with cervical cytology report of atypical squamous cells and the diagnostic role of high-risk HPV infection. *European Journal of Gynaecological Oncology*. 2013;34:159-162.
415. Yuan X, Yang Y, Gu D et al. Prevalence of human papillomavirus infection among women with and without normal cervical histology in Shandong Province, China. *Archives of Gynecology & Obstetrics*. 2011;283:1385-1389.
416. Zappacosta R, Caraceni D, Ciccocioppo L et al. Implementing specificity of HPV-DNA primary screening in a successful organised cervical cancer prevention programme. *Gynecol Oncol*. 2013;128:427-432.
417. Zeferino LC, Rabelo-Santos SH, Villa LL et al. Value of HPV-DNA test in women with cytological diagnosis of atypical glandular cells (AGC). *European Journal of Obstetrics, Gynecology, & Reproductive Biology*. 2011;159:160-164.
418. Zeng WJ, Li Y, Fei HL et al. The value of p16ink4a expression by fluorescence in situ hybridization in triage for high risk HPV positive in cervical cancer screening. *Gynecol Oncol*. 2011;120:84-88.
419. Zhang R, Velicer C, Chen W et al. Human papillomavirus genotype distribution in cervical intraepithelial neoplasia grades 1 or worse among 4215 Chinese women in a population-based study. *Cancer Epidemiology*. 2013;37:939-945.
420. Zhang R, He YF, Chen M et al. Diagnosis of 25 genotypes of human papillomaviruses for their physical statuses in cervical precancerous/cancerous lesions: a comparison of E2/E6E7 ratio-based vs. multiple E1-L1/E6E7 ratio-based detection techniques. *Journal of Translational Medicine*. 2014;12:282.
421. Zhao FH, Jeronimo J, Qiao YL et al. An evaluation of novel, lower-cost molecular screening tests for human papillomavirus in rural China. *Cancer Prev Res*. 2013;6:938-948.
422. Zorzi M, Del MA, Farruggio A et al. Use of a high-risk human papillomavirus DNA test as the primary test in a cervical cancer screening programme: a population-based cohort study. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2013;120:1260-1267.