

**READ SAFETY DIRECTIONS  
BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY**



# CLiKziN™

## SPRAY-ON


### SHEEP BLOWFLY TREATMENT

Active Constituent: 12.5 g/L DICYCLANIL

**READ DIRECTIONS  
FOR USE BEFORE  
OPENING OR USING  
THIS PRODUCT**

Elanco Australasia Pty Ltd  
ABN 64 076 745 198  
112 Wharf Road, West Ryde  
NSW 2114, Australia

Elanco, CLiKziN and the diagonal bar are  
trademarks owned or licensed by  
Eli Lilly and Company, its subsidiaries or affiliates.



For the protection of sheep  
against fly strike (*Lucilia  
cuprina*) for up to 11 weeks.  
For the protection of marking  
wounds on lambs against fly  
strike (*Lucilia cuprina*) for up  
to 11 weeks.

**20 Litres**



YL400202ALA 916800

# CLiKziN™

## SPRAY-ON

### GENERAL INSTRUCTIONS

CLiKziN provides protection to sheep against body, breech and poll strike for up to 11 weeks and may be used on sheep with any length wool from 3 weeks after shearing. CLiKziN also protects marking wounds on lambs against fly strike for up to 11 weeks.

The dose volume is determined by the weight of sheep being treated. Apply CLiKziN according to the diagrams and tables in the Directions for Use.

**DO NOT DILUTE.** Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by Elanco. CLiKziN may be applied to all breeds of sheep to obtain protection against fly strike. CLiKziN is applied as a low-volume treatment onto areas where protection is required. When applying CLiKziN, the nozzle of the applicator gun should be high enough above the fleece (approximately 15–25 cm), to achieve a 15 cm wide band at each pass for protection from body strike.

**DO NOT** apply during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for marking to be carried out at times of minimal fly activity.

Reduced efficacy may occur in sheep affected with dermatophilosis (dermo/lumpy wool).

Significant soiling with dags, mud, dust or urine may reduce efficacy.

Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.

### SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product wear elbow-length rubber gloves. Wash hands after use.

### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

### ADDITIONAL USER SAFETY INFORMATION

**Sheep Rehhandling Interval:** After treatment wait until sheep are dry before rehhandling.

If additional hazard information is required, refer to the Safety Data Sheet. For a copy phone 1800 226 324.

### ENVIRONMENTAL PROTECTION

DO NOT contaminate dams, rivers or streams with the product or used containers.

### DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinseate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

### STORAGE

Keep out of reach of children. Store below 30°C (Room Temperature). Protect from frost. Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

### EXCLUSION OF LIABILITY

To the extent permitted by legislation, Elanco Australasia Pty Ltd will not be liable or responsible for any loss, damage or expense where the product is not used as directed.

### DIRECTIONS FOR USE

#### Restraints:

**DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption.**

**DO NOT USE on sheep off-shears.**

#### Contraindications:

**DO NOT dilute.**

**DO NOT mix with any other product.**

#### FOR THE PREVENTION OF BODY STRIKE (Diagram 1)

SHEEP WEIGHT (kg)	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER 20 L
10-20	10	10	1	2000
21-30	12	12	1	1666
31-50	15	7.5	2	1333
>50	18	9	2	1111

#### FOR THE PREVENTION OF BREECH STRIKE (Diagram 1)

SHEEP WEIGHT (kg)	TOTAL VOLUME/SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER 20 L
10-20	10	10	1	2000
21-30	12	12	1	1666
31-50	15	7.5	2	1333
>50	18	9	2	1111

#### FOR THE PREVENTION OF POLL STRIKE

Rams and sheep susceptible to poll strike should be treated. Apply 5 mL of CLiKziN to the areas requiring protection, around the poll and horns.

#### FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (Diagram 2)

- When a tail has been amputated, apply 5 mL over the crutch ensuring sufficient coverage of the open wound.
- When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied.
- In the case of ram lambs a further 5 mL can be sprayed over and around the scrotum area after the rubber ring has been applied or after knife castration.

LAMBS WEIGHT (< 30 kg)	TOTAL VOLUME/LAMB (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	AVERAGE NO. OF LAMBS TREATED PER 20 L
Ram lambs	10	5	2	2666
Ewe lambs	5	5	1	

#### CRITICAL COMMENTS

1. Use only a spray-on applicator gun recommended by Elanco.
2. Ensure the applicator gun is clean.
3. Calibrate applicator gun using an accurate volume measure.
4. Ensure all areas to be protected are treated.
5. When preventing body strike ensure bands overlap along the midline – see Diagram 1.
6. When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool – see Diagram 2.
7. This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes.
8. Heavy rain following the application of this product could diminish the period of protection.
9. CLiKziN contains a scourable colouring agent for easy band identification.
10. Withholding periods must be observed following application of CLiKziN for any purpose including after marking as finite residues will occur.

Diagram 1




Diagram 2



**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 7 days before slaughter for human consumption.**

**MILK: DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption.**

**WOOL HARVEST INTERVAL: DO NOT USE less than 1 month before shearing or fibre collection.**

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 226 324 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

APVMA Approval No: 64628/100213

**☎ CUSTOMER INFORMATION**

Australia: 1800 226 324 TOLL FREE

Batch No.	
Expiry Date	



YL400202ALB 916800