



Product Name: FASINEX 240 ORAL FLUKICIDE FOR CATTLE AND SHEEP
APVMA Approval No: 51262/115927

Label Name:	FASINEX 240 ORAL FLUKICIDE FOR CATTLE AND SHEEP
-------------	---

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	Active Constituent: 240g/L TRICLABENDAZOLE
-------------------------	--

Claims:	FASINEX 240 contains triclabendazole, a member of the benzimidazole family of drenches. It is effective against triclabendazole sensitive strains of <i>Fasciola hepatica</i> (liver fluke). For the treatment of triclabendazole-susceptible Early Immature, Immature and Mature liver fluke (<i>Fasciola hepatica</i>) in cattle and sheep. Resistance may develop to any drench.
---------	---

Net Contents:	2.2 Litres, 5 Litres, 13 Litres
---------------	---------------------------------

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE in lactating animals where milk or milk products may be used or processed for human consumption.
------------	---

Precautions:	In areas where Black Disease is known to occur, adequate vaccination procedures are advised.
--------------	--

Dosage and Administration:	SHAKE WELL BEFORE USE See attached dose tables for Cattle and Sheep. CATTLE CRITICAL COMMENTS: 1. FASINEX 240 may be administered at the same time as animals are treated for lice.
----------------------------	--

2. FASINEX 240 may be administered to calves, pregnant cows, or weak or debilitated animals.
 3. Wash out equipment with water after use.
 4. FASINEX 240 is palatable to cattle.
 5. Cattle in excess of 650kg are to be dosed at a rate of 2.5mL per 50kg bodyweight. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves).
- DO NOT underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

SHEEP CRITICAL COMMENTS

1. FASINEX 240 may be administered to lambs, pregnant or lactating ewes, or weak or debilitated animals.
2. FASINEX 240 does not stain wool.
3. Wash out equipment with water after use.
4. FASINEX 240 can be used at the same time as jetting with other ectoparasiticides.
5. FASINEX 240 is palatable to sheep.
6. Sheep in excess of 75kg should be dosed at a rate of 0.5mL per 10kg bodyweight. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, lambs, wethers, rams). Where there is large variation in size within the group, dose rate should be based on the label directions for each weight range. DO NOT underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

This section contains file attachment.

File Name: Fasinex 240 Proposed Dose Table Final.docx

File Size: 15554 bytes

General Directions:

FASINEX 240 Oral Flukicide for Cattle and Sheep is a ready to use, free flowing aqueous suspension for the treatment and control of acute and chronic fascioliasis of cattle and sheep.

FASINEX 240 when used at the recommended rate of 12mg triclabendazole/kg bodyweight for cattle (i.e. 2.5ml/50kg BW) and a minimum dose of 10mg triclabendazole/kg bodyweight for sheep, is highly effective against susceptible early immature fluke as well as immature and adult fluke, and has a high safety margin. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.

Withholding Periods:

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 21 days before slaughter for human consumption

MILK: DO NOT USE in lactating animals where milk or milk products may be used or processed for human consumption. DO NOT USE less than 21 days before calving or lambing in cows or ewes where milk and milk products from treated animals may be used for human consumption.

Where FASINEX 240 is accidentally given within this period or should any cows or ewes calve or lamb earlier than 21 days after treatment, milk will contain residues. This milk must not be used for human consumption, or supplied for processing for at least 21 days following treatment. Calves fed this milk should not be slaughtered for human consumption within 10 days. Lambs fed this milk should not be slaughtered for human consumption within 7 days.

Trade Advice:

EXPORT SLAUGHTER INTERVAL (ESI):

(CATTLE) DO NOT USE less than 56 days before slaughter for export.

(SHEEP) DO NOT USE less than 63 days before slaughter for export.

The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on 1800 226 324 or the APVMA website (www.apvma.gov.au/residues).

Safety Directions: Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

First Aid Instructions: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

First Aid Warnings:

Additional User Safety: Additional information is listed in the safety data sheet.

Environmental Statements: ENVIRONMENTAL PROTECTION
Dangerous to fish. DO NOT contaminate dams, rivers or streams with the drench or used container.

Disposal: Disposal: This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Storage: Store below 30 deg C (Room temperature) in original container, tightly closed in a cool, dry place. Protect from frost.

FASINEX 240 Proposed Dosing Table for Cattle and Sheep

Cattle BodyWeight (kg)	Oral Dose (mL)
Up to 100 kg	0.5 mL/10 kg liveweight
101-150 kg	7.5
151-200 kg	10
201-250 kg	12.5
251-300 kg	15
301-400 kg	20
401-500 kg	25
501-600 kg	30
601-650 kg	32.5
Over 650 kg	2.5 mL/50 kg liveweight

Sheep BodyWeight (kg)	Oral Dose (mL)
15-20 kg	1.0
21-30 kg	1.5
31-40 kg	2.0
41-50 kg	2.5
51-60 kg	3.0
61-70 kg	3.5
Over 70kg	0.5ml/10 kg liveweight