

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

Elanco™

AF0602

Paylean™ 20

TM

10kg NET

Ractopamine Hydrochloride Premix

Active constituent: 20g/kg ractopamine hydrochloride (equivalent to 17.8 g/kg ractopamine)

Pigs: Finisher pigs: For increased rate of weight gain, improved feed efficiency, increased carcass leanness and increased dressing percentage.

Sows: For reduced lactation weight loss in breeding sows.

Turkeys: For increased rate of weight gain and improved feed efficiency in finisher turkeys.

DIRECTIONS FOR USE:

Restraint:

DO NOT ADMINISTER PAYLEAN TO SPECIES OTHER THAN PIGS AND TURKEYS.

DO NOT USE in turkeys which are producing or may in the future produce eggs for human consumption.

DO NOT USE IN BREEDING TURKEYS.

FOR USE IN PIG AND FINIHSER TURKEY FEEDS ONLY.

Note to feed formulators:

Unintended inclusion of medication into animal feeds may result in residues that adversely impact trade access or the rules of professional competition. Use cleaning or flushing, or avoid sequencing feed for other species (in particular ruminant and horse feeds) immediately after manufacture of pig or turkey finisher feeds containing Paylean.

A copy of directions for use relevant to final feed are required to be provided to the producers with each consignment of Paylean treated feed. Contact Elanco Animal Health for further information.

Precautions:

Pigs fed Paylean are at an increased risk for exhibiting the downer pig syndrome (could also be referred to as leg weakness syndrome). Pig handling methods to reduce the incidence of downer pigs should be thoroughly evaluated prior to initiating use of Paylean. Feeding ractopamine to tom turkeys during periods of excessive heat can result in increased mortality.

Dosage and Administration:

Pigs

Recommended dosages of ractopamine hydrochloride:

- 5-20 ppm in final feed for increased rate of weight gain and improved feed efficiency.
- 10-20 ppm in final feed for increased carcass leanness and increased carcass dressing percent.
- 20 ppm in lactating sows to reduce weight loss.

Pigs Mixing Directions:

PAYLEAN 20 PREMIX MUST BE THOROUGHLY MIXED INTO FEEDS BEFORE USE.

All calculations are based on 90% dry matter.

Ractopamine Concentration in Final Feed g/tonne (ppm)	Grams of Paylean 20 Premix to Add per Tonne of Final Feed
5	250
10	500
15	750
20	1000 (1 kg)

IMPORTANT – To ensure adequate mixing, it is recommended to pre-blend Paylean 20 Premix in a small quantity of feed before incorporation into the total amount of complete feed. Thoroughly mix the pre-blend into a nutritionally balanced pig ration to result in a final ration containing 5 to 20 ppm of ractopamine. Amino acid content of the ration may influence response to treatment. Dietary specifications should be determined in consultation with a recognised pig nutritionist in order to optimise Paylean benefits.

Feeding directions: Feed complete ration containing ractopamine continuously as the sole ration to finisher pigs for the last 28-42 days prior to slaughter.

To achieve best results add 20 ppm ractopamine to sow diets throughout lactation.

Turkeys

The recommended dose for finisher hen and tom turkeys is 5 – 13 ppm for increased rate of weight gain and improved feed efficiency.

Turkeys Mixing Directions:

PAYLEAN 20 PREMIX MUST BE THOROUGHLY MIXED INTO FEEDS BEFORE USE.

Ractopamine Concentration in Final Feed g/tonne (ppm)	Grams of Paylean 20 Premix to Add per Tonne of Final Feed
5	250
9	450
13	650

IMPORTANT – To ensure adequate mixing, it is recommended to pre-blend Paylean 20 Premix in a small quantity of feed before incorporation into the total amount of complete feed. Thoroughly mix the pre-blend into a nutritionally balanced turkey ration to result in a final ration containing 5 to 13 ppm ractopamine.

Feeding directions: Feed complete ration containing ractopamine continuously as the sole ration to finisher turkeys for the last 14 days prior to slaughter.

WARNING: Not for use in humans. Individuals with cardiovascular disease should exercise special caution to avoid exposure. The active ingredient in Paylean, ractopamine hydrochloride, is a beta-adrenergic agonist. Additional information is in the Safety Data Sheet.

WITHHOLDING PERIODS:

PIGS: MEAT - DO NOT USE less than 12 hours before slaughter for human consumption.

TURKEYS: REMOVE ALL MEDICATED FEED 6 hours before slaughter for human consumption.

EGGS: DO NOT USE in turkey hens which are producing or may in the future produce eggs for human consumption or processing.

EXPORT TRADE ADVICE – TREATED STOCK: Exporters need to comply with the regulations/standards of importing countries with regard to the use of animal health products in livestock. Producers are advised to contact their export slaughter facility for information before giving pigs or turkeys feed to which this product has been added.

SAFETY DIRECTIONS: Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. If product in eyes, wash it out immediately with water. Wash hands after use. When opening the container and preparing stock feed wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat, elbow length PVC gloves and face shield or goggles. After each day's use, wash gloves and contaminated clothing.

FIRST AID INSTRUCTIONS: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126.

DISPOSAL OF BAGS: Single-rinse or shake container into the medicated feed. Do not dispose of undiluted chemicals on site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Additional information is in the Material Safety Data Sheet.

Manufactured in U.K. for:
Elanco Australasia Pty Ltd.
112 Wharf Road, West Ryde, NSW. 2114
TEL: Toll free 1800 226 324

APVMA Approval Number: 54172/109774

STORAGE: Store below 30°C (Room temperature) in a dry place.

For batch number and expiry date see reverse.

USE NO HOOKS



Paylean, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

Avoid breathing dust. In case of inadequate ventilation wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace.
IF INHALED: if breathing is difficult, remove victim to fresh air and keep at rest in a position comfortable for breathing.

Lot

DOM






EXP

YL209745X

PRODUCT INFO

BLUE #:	209745A	Product Name:	Paylean 20
Item Code:	YL209745X	Component:	POD Label
Product Code:	AF0602	Pack Size:	10kg
Previous Item Code	YL401893X		

ARTWORK INFO

Template:	LA_209x500mm_10kg_25kg_POD_v1	Packaging Spec(s):	Primary Spec: B6G090 Base Spec: CM2949		
Barcodes/Type:	EAN 13: 9323829002736	Add. Info:	PMS 348 C converted to CMYK for digital printing		
		Minimum Core Data Point Size:	8 pt		
GTIN:	N/A	Proof #	P2b	By/Date	DM 16-Jun-2018
					
NON PRINTING	CYAN	MAGENTA	YELLOW	BLACK	

ELANCO ARTWORK LEGEND v13