CAUTION

KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING FOR ANIMAL TREATMENT ONLY



CREDELIO 56.25 mg CHEWABLE TABLETS FOR DOGS Each chewable tablet contains 56.25 mg Lotilaner CREDELIO 112.5 mg CHEWABLE TABLETS FOR DOGS Each chewable tablet contains 112.5 mg Lotilaner CREDELIO 225 mg CHEWABLE TABLETS FOR DOGS Each chewable tablet contains 225 mg Lotilaner CREDELIO 450 mg CHEWABLE TABLETS FOR DOGS Each chewable tablet contains 450 mg Lotilaner CREDELIO 900 mg CHEWABLE TABLETS FOR DOGS

Each chewable tablet contains 900 mg Lotilaner

Indications

Credelio provides immediate treatment and persistent killing activity for up to 1 month for fleas (Ctenocephalides felis). Rapidly kills fleas before they have a chance to lay eggs. Controls flea allergy dermatitis (FAD) by controlling adult fleas. For the treatment and control of Paralysis Ticks (Ixodes holocyclus), Brown Dog Ticks (Rhipicephalus sanguineus) and Bush Ticks (Haemaphysalis longicornis) for one month.

Ixodes holocyclus ticks do not occur in Western Australia. In tick season, daily searching for, and removal of, any ticks found is recommended.

DIRECTIONS FOR USE

The safe use of CREDELIO in breeding males and females has not been evaluated. Not recommended for use on puppies under 8 weeks of age or weighing less than 1.3 kg.

Dosage and Administration

CREDELIO is available in five strengths.

The recommended dosage for 1.3 to 2.5 kg dogs is one CREDELIO 56.25 mg Chewable Tablet.

The recommended dosage for > 2.5 to 5.5 kg dogs is one CREDELIO 112.5 mg Chewable Tablet.

The recommended dosage for > 5.5 to 11 kg dogs is one CREDELIO 225 mg Chewable Tablet.

The recommended dosage for > 11 to 22 kg dogs is one CREDELIO 450 mg Chewable Tablet.

The recommended dosage for > 22 to 45 kg dogs is one CREDELIO 900 mg Chewable Tablet.

The minimum recommended dose rate is 20 mg Lotilaner/kg bodyweight administered with or following food. Dogs should be dosed appropriately to ensure a dose rate of 20 to 43 mg Lotilaner/kg bodyweight.

Recommended Dose Schedule.

| Dog Weight | CREDELIO Chewable Tablet Strength | Tablets Administered |
|----------------|-----------------------------------|----------------------|
| 1.3 – 2.5 kg | 56.25 mg | One |
| > 2.5 – 5.5 kg | 112.5 mg | One |
| > 5.5 – 11 kg | 225 mg | One |
| > 11 – 22 kg | 450 mg | One |
| > 22 - 45 kg | 900 mg | One |

For dogs weighing above 45 kg bodyweight, the appropriate combination of tablets matching the weight of the dog should be used. For dogs weighing between 1.3 to 45 kg, an appropriate combination of tablets may also be used, provided the minimum dose rate of 20 mg/kg is maintained. Within each weight range, and for any suitable combination of tablets, whole tablets must be used. Administer CREDELIO with or following food.

Treat your dog on or about the same day each month. CREDELIO may be used all year round.

General Directions

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

CREDELIO was shown to be safe in various breeds of dogs and in puppies 8 weeks of age and older and in dogs weighing 1.3 kg and above. No adverse reactions were observed following oral administration to puppies at 8 weeks of age and weighing 1.3 - 3.6 kg treated with overdoses of up to 5 times the maximum recommended dose on eight occasions at monthly intervals.

Lotilaner is part of the isoxazoline family of chemicals. Adverse reactions to this family of chemicals are rarely observed but may include vomiting, diarrhoea, lethargy, inappetance, itching and very rarely, seizures. Most adverse reactions are self-limiting and of short duration. If you have any concerns, please speak to your veterinarian.

General Instructions

As CREDELIO is an oral treatment, there are no restrictions on swimming, bathing and shampooing before or after administration. CREDELIO contains Lotilaner, a member of the isoxazoline antiparasitic class. Lotilaner acts antagonistically on arthropod ligand-gated chloride

channels (GABA-receptor and glutamate receptor).

CREDELIO starts to kill fleas within 2 hours from administration, with fleas being eliminated within 6-12 hours. For new infestations, fleas are eliminated within 4-6 hours throughout the entire month. CREDELIO kills existing and newly emerged fleas on dogs before they can lay eggs due to its rapid onset of action and long lasting efficacy. Therefore, CREDELIO breaks the flea life cycle and prevents environmental contamination in areas

CREDELIO can be used as part of a treatment strategy for flea allergy dermatitis (FAD).

CREDELIO demonstrated > 95% efficacy for a full month against Paralysis Ticks, Brown Dog Ticks and Bush Ticks.

CREDELIO is suitable for use in puppies from 8 weeks of age and a minimum bodyweight of 1.3 kg.

SAFETY DIRECTIONS

May irritate eyes. Avoid contact with eyes. Wash hands after use.

If poisoning occurs, contact a doctor or Poisons Information Centre. (Phone Australia 131 126).

ADDITIONAL USER SAFETY INFORMATION

Safety Data Sheet: If additional hazard information is required, refer to the Safety Data Sheet. For a copy phone 1800 226 324.

Dispose of container by wrapping with paper and putting in garbage.

STORAGE

Store below 30 °C (room temperature)

Elanco Australasia Pty Ltd 112 Wharf Road

West Ryde NSW 2114 Australia Tel: 1800 995 709

APVMA Approval No.: 82853/106969; 82854/106970; 82855/106971; 82856/106972; 82857/106973

Presentation

1 tablet, 3 tablets or 6 tablets per carton.

Exclusion of Liability

To the extent permitted by legislation, Elanco Australasia Pty Ltd will not be liable or responsible for any loss, damage or expense where the product is not used as directed.

Credelio, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.





