

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY



## FASIMEC™ CATTLE

Active Constituents:  
120 g/L TRICLABENDAZOLE  
2 g/L IVERMECTIN

SHAKE WELL BEFORE USE

15 Litres



Elanco™

### Oral Flukicide and Broad Spectrum Drench

For the treatment of triclabendazole-sensitive Early Immature, Immature and Mature liver fluke (*Fasciola hepatica*) in cattle.

For the treatment and control of ivermectin-sensitive strains of roundworms and lungworm (including benzimidazole, levamisole, and morantel-resistant strains), and sucking lice in cattle.

YL400091ALA

## Parasites treated and controlled by Fasimec Cattle

Early immature, immature and adult liver fluke ..... *Fasciola hepatica*

### Adult and immature (L4) stages of:

Small brown stomach worm (including inhibited stages) ..... *Ostertagia ostertagi*  
Barber's pole worm ..... *Haemonchus placei*  
Stomach hair worm ..... *Trichostrongylus axei*  
Intestinal hair worm ..... *Trichostrongylus colubriformis*  
Small intestinal worm ..... *Cooperia* spp.  
Nodule worm ..... *Oesophagostomum radiatum*

### Adult stages of:

Large bowel worm ..... *Oesophagostomum venulosum*  
Whipworm ..... *Trichuris* spp.  
Intestinal threadworm ..... *Strongyloides papillosus*

### Immature stages of:

Thin necked intestinal worm ..... *Nematodirus* spp.

Lungworm, Adult and immature ..... *Dictyocaulus viviparus*

### Sucking lice:




Long-nosed sucking louse ..... *Linognathus vituli*  
Short-nosed sucking louse ..... *Haematopinus eurysternus*  
Small blue sucking louse ..... *Solenopotes capillatus*  
Does not provide control of the cattle biting louse, *Bovicola (Damalinia) bovis*.

## DIRECTIONS FOR USE

**Restraints: NOT TO BE USED FOR ANIMALS PRODUCING MILK FOR HUMAN CONSUMPTION OR PROCESSING.**

### SHAKE WELL BEFORE USE

**ORAL DOSE RATE: CATTLE 5 mL per 50 kg BW**

Parasites Controlled	 Cattle weight	 Oral Dose (mL)	 Number treated per pack
Liver fluke,	up to 100 kg	1 mL/10 kg bodyweight	
Gastrointestinal	101-150 kg	15	1000
roundworms, Lungworm	151-200 kg	20	750
and Sucking Lice	201-250 kg	25	600
	251-300 kg	30	500
	301-400 kg	40	375
	401-500 kg	50	300
	501-600 kg	60	250
	601-650 kg	65	230
	Over 650 kg	5 mL/50 kg bodyweight	

**Critical Comments**

1. FASIMEC may be administered at the same time as animals are treated for biting lice.
2. FASIMEC may be administered to calves, pregnant cows or weak or debilitated animals.
3. Wash out equipment with water after use.
4. FASIMEC is palatable to cattle.
5. Cattle in excess of 650 kg should be dosed at 5 mL per 50 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). DO NOT underdose. Where there is a larger variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS

**MEAT - DO NOT USE less than 21 days before slaughter for human consumption.**

**MILK - DO NOT USE in lactating cows or within 28 days of calving where milk or milk products may be used for human consumption.**

### TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 56 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on (1800 226 324) or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**

**Protection of Wildlife, Fish, Crustaceans and the Environment:** Ivermectin is extremely dangerous to aquatic species. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used containers.

### GENERAL INSTRUCTIONS:

#### SHAKE WELL BEFORE USE

**Precaution:** In areas where Black Disease is known to occur, adequate vaccination procedures are advised.

FASIMEC Cattle is a ready to use, free flowing aqueous suspension for the treatment and control of acute and chronic fascioliasis, mixed infections of gastrointestinal roundworms, lungworm and sucking lice.

Fasimec Cattle contains triclabendazole, a member of the benzimidazole family of drenches, and ivermectin, a member of the avermectin (*macrocyclic lactone*) family of drenches.

Fasimec Cattle when used at the recommended rate of 12 mg triclabendazole and 0.2 mg ivermectin per kg bodyweight for cattle, i.e. 5 mL Fasimec Cattle per 50 kg bodyweight, combines high efficacy against all ages of triclabendazole-sensitive liver fluke, ivermectin-sensitive adult and immature roundworms, lungworm and sucking lice, with a high safety margin. Resistance may develop to any drench.

Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.

**SAFETY DIRECTIONS: May irritate the eyes. Avoid contact with eyes. Wash hands after use.**

**FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre (Ph: 13 1126).**

**Material Safety Data Sheet:** If additional hazard information is required refer to the Material Safety Data Sheet. For a copy phone 1800 226 324.

**Disposal:** This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

**Storage:** Store below 30°C (Room Temperature) in original container, tightly closed, in a cool, dry place. Protect from light.

### Exclusion of liability

To the extent permitted by legislation, Elanco Australasia Pty Ltd will not be liable or responsible for any loss, damage or expense where the product is not used as directed.

APVMA Approval No. 52406/15L/0109

Elanco, Fasimec and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Batch Number	
Expiry Date	



YL400091ALB



Elanco Australasia Pty Ltd, ABN 64 076 745 198, 112 Wharf Road, West Ryde, NSW 2114, Australia

 CUSTOMER INFORMATION Australia: 1800 226 324 TOLL FREE