DIRECTIONS FOR USE

Restraint

DO NOT USE in dairy animals producing or which will in the future be producing milk for human consumption.
This product is contraindicated for use in calves under 50kg bodyweight.

Dosage and Administration: FASIMEC CATTLE Pour-On dose rate is 1 mL / 10 kg bodyweight. Apply along the backline of the animal, using a standard pour-on applicator, in a strip starting between the shoulder blades. This product is not to be given orally or injected.

Liveweight (kg)	Dose Volume (mL)	Doses per pack 15 L
51-75	7.5	2000
76-100	10	1500
101-125	12.5	1200
126-150	15	1000
151-175	17.5	857
176-200	20	750
201-250	25	600
251-300	30	500
301-350	35	428
351-400	40	375
401-450	45	333
451-500	50	300
501-550	55	272
551-600	60	250
601-650	65	230

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

For cattle over 650 kg, dose at rate of 1 mL / 10 kg bodyweight. A representative sample of animals should be weighed before treatment either with scales or weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). DO NOT underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

FASIMEC CATTLE Pour-On is rainfast. Rain occurring as early as 2 hours after treatment will not affect the efficacy of the product.

A white deposit may form on the back of treated animals. This is quite normal. It is not a sign of treatment failure and should not be interfered with. DO NOT brush off or wash off. This deposit may last for a few days or weeks depending on weather conditions. Temporary skin reaction may occur along the application site, especially if applied during very hot weather. This should resolve without treatment. If signs persist or worsen, contact your veterinarian. After use clean all equipment with cooking oil. DO NOT use water.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 49 days before slaughter for human consumption. Beef calves born to cows that were treated with FASIMEC CATTLE POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for human consumption for 49 days after treatment of the dam.

MILK: DO NOT USE in dairy animals producing or which will in the future be producing milk for human consumption.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 140 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Elanco Australasia Pty Ltd on (1800 226 324) or the APVMA website (www.apvma.gov.au/residues).

Beef calves born to cows that were treated with FASIMEC CATTLE POUR-ON during pregnancy and calves suckling cows that have been treated during lactation must not be slaughtered for the export market for 140 days after treatment of the dam.

SAFETY DIRECTIONS: Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. If product in eyes, first quickly, gently and completely blot or wipe chemical from eyes then flush with water. Wash hands after use. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre, Phone 13 1126. If skin contact occurs, remove contaminated clothing quickly and gently blot or wipe away chemical with a tissue or cloth, preferably impregnated with cooking oil, then wash skin thoroughly using soap and water. DO NOT use water first.

Material Safety Data Sheet:
If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy phone 1800 226 324.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT:
Abamectin is extremely toxic to aquatic species. DO NOT contaminate dams, rivers, streams or waterways with the chemical or used containers.

DISPOSAL:
This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo. Triple rinse containers before disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. DO NOT burn empty containers or product.

STORAGE: Store below 30°C (Room Temperature).

APVMA Approval No. 58971/15L/0609

Elanco Australasia Pty Ltd, ABN 64 076 745 198, 112 Wharf Road, West Ryde, NSW 2114, Australia
☎ CUSTOMER INFORMATION Australia: 1800 226 324 TOLL FREE

Elanco™

YL400132ALB

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

FASIMEC™ CATTLE POUR-ON

Active Constituents:
5 mg/mL ABAMECTIN
300 mg/mL TRICLABENDAZOLE

Flukicide and Broad Spectrum Anthelmintic

For the treatment and control of roundworms, liver fluke (all 3 stages) and external parasites of beef cattle.

15 Litres

Elanco, Fasimec and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

Elanco™

YL400132ALA