



Product Name: Elanco Flubenol Oral Wormer for Pigs and Chickens  
APVMA Approval No: 86633/116234

Label Name:	Elanco Flubenol Oral Wormer for Pigs and Chickens
-------------	---

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	Flubendazole 50 mg/g
-------------------------	----------------------

Claims:	Flubenol contains flubendazole, a member of the benzimidazole group family of chemicals. Flubenol can be used as an aid for the control of sensitive strains of the following internal parasites: Gastrointestinal worms ( <i>Ascaridia galli</i> , <i>Heterakis gallinarum</i> , <i>Capillaria</i> spp., <i>Railletina</i> spp.) in chickens and Gastrointestinal worms ( <i>Ascaris suum</i> , <i>Hyostrogylus rubidis</i> , <i>Oesophagostomum dentatum</i> ) and lungworm ( <i>Metastrongylus apri</i> ) in pigs. Resistance may develop to any chemical
---------	---

Net Contents:	600 g 2 kg 12 kg 25 kg
---------------	---------------------------------

Directions for Use:	
---------------------	--

Restrains:	
------------	--

Contraindications:	This product is contraindicated for use in pigeons and parrots.
--------------------	---

Precautions:	Do not overdose
--------------	-----------------

Side Effects:	
---------------	--

Dosage and Administration:	<p><b>DOSAGE and ADMINISTRATION</b></p> <p>Use treated feeds within 6 months of preparation.</p> <p><b>Chickens</b>  <i>Ascaridia galli</i>, <i>Heterakis gallinarum</i>, <i>Capillaria</i> spp:</p> <p>600 g of Flubenol per tonne of feed (equivalent to 30 g flubendazole (30 ppm)) for 7 consecutive days.</p> <p>For small numbers of animals, 1 supplied measuring spoon (13g) of Flubenol is mixed into 21.5 kg of compound feed for chickens.</p> <p><b>Tapeworm (<i>Raillietina</i> spp.):</b>  1200 g of Flubenol per tonne of feed (equivalent to 60 g flubendazole (60 ppm)) for 7 consecutive days.</p> <p>Ensure birds have access to feed to enable consumption of the required dose. Optimum results are achieved when using stringent chicken housing hygiene requirements.</p> <p><b>Pigs</b>  <i>Ascaris suum</i>, <i>Hyostrogylus rubidis</i>, <i>Oesophagostomum dentatum</i> and <i>Metastrongylus apri</i></p> <p>600 g of Flubenol per tonne of feed (equivalent to 30 g flubendazole (30 ppm)) for 10 consecutive days.</p> <p>Individual and single administration to breeding pigs:  Flubenol is mixed into the feed at a dose of 5 mg flubendazole per kg body weight. This corresponds to one supplied measuring spoon (13 g) of Flubenol per 130 kg of body weight.</p> <p>Flubenol should be thoroughly mixed into feed to ensure even distribution of the medication. Feed can be administered as either a mash, crumble or pellet.</p>
----------------------------	---

General Directions:	
---------------------	--

Withholding Periods:	<p><b>MEAT (CHICKEN): REMOVE ALL MEDICATED FEED 7 days before slaughter for human consumption.</b></p> <p><b>EGGS: Zero (0) Days</b></p> <p><b>MEAT (PIG): REMOVE ALL MEDICATED FEED 7 days before slaughter for human consumption.</b></p>
----------------------	---

Trade Advice:	<p><b>TRADE ADVICE</b></p> <p><b>PIGS: (ESI) DO NOT USE less than 7 days before slaughter for export. Before using this product, confirm the current ESI from Elanco Australasia on 1800226324 or the APVMA website (<a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a>)</b></p>
---------------	--

Safety Directions:	<p>May irritate the eyes. Avoid contact with eyes. When opening the container and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.
-------------------------	--

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	<b>ENVIRONMENTAL PROTECTION.</b> Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.
---------------------------	---

Disposal:	600g container: Dispose of container by wrapping with paper and putting in garbage.  2-25 kg bags: Single-rinse or shake container into the medicated feed. Do not dispose of undiluted chemicals on-site. Puncture bag and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the container 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------	--

Storage:	Store below 30 degrees Celsius (Room temperature). Use treated feeds within 6 months of preparation.
----------	--