

DIRECTIONS FOR USE

Restraints

DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption.

Remove wool from over the wound using clippers or shears. Apply directly on to the wound and 25 mm into the surrounding wool until area is wet and 'blue'. (A typical clipped flystrike wound, 200 cm² in area, should take approximately 6 seconds to treat).

SHAKE WELL BEFORE USE.

NOT TO BE USED FOR ANY OTHER PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: NIL

MILK: DO NOT USE on ewes which are producing or may in the future produce milk or milk products for human consumption.

WOOL: NIL

An export slaughter interval (ESI) may apply to this product. Contact Elanco Animal Health before using on sheep destined for the export market.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

SAFETY DIRECTIONS: Will irritate the eyes and skin.

Avoid contact with eyes and skin. Wash hands after use.

When using the prepared spray wear elbow-length PVC gloves.

After each day's use, wash gloves.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13 1126. If sprayed on skin, wash thoroughly. If sprayed in mouth, rinse mouth with water.

ADDITIONAL USER SAFETY INFORMATION

Sheep Rehandling Interval: After treatment wait until sheep are dry before rehandling.

DISPOSAL: Aerosols should be completely emptied before disposal. Place empty can in household rubbish. Do not puncture or incinerate.

PROPELLANT: Hydrocarbon

Elanco Animal Health
A Division Of Eli Lilly Australia Pty. Limited
A.B.N. 39 000 233 992
112 Wharf Road, West Ryde, N.S.W. 2114
Tel: Toll free 1800 226 324

APVMA Approval Number 56734/0614

STORAGE: Store below 30° C
(Room temperature).


KEEP OUT OF REACH OF CHILDREN

For batch and expiry see base of container.

Elanco, Extinosad and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

PRESSURISED DISPENSER, PROTECT FROM SUNLIGHT AND DO NOT EXPOSE TO TEMPERATURES EXCEEDING 50° C. DO NOT PIERCE OR BURN, EVEN AFTER USE.

EXTREMELY FLAMMABLE. KEEP AWAY FROM NAKED FLAME OR ANY INCANDESCENT MATERIAL. KEEP AWAY FROM SOURCES OF IGNITION-NO SMOKING.



Aerosol for Wounds

370 g NET

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

Elanco™

AH0498



Extinosad™

TM



AEROSOL FOR WOUNDS

ACTIVE CONSTITUENTS: 2.80 g/kg spinosad
0.39 g/kg chlorhexidine digluconate minimum



For the treatment and prevention of blowfly (*Lucilia cuprina*) strike in mulesing, marking and other wounds of sheep, including strains resistant to organophosphates. Contains an antiseptic.

YL089820ALX